TAXONOMIC INDEX

Copeia 2011, Nos. 1–4

NEW NAMES

Congoglanis 477–489
inga 477–489
sagitta 477–489
Creteuchiloglanis 226–241
brachypterus 226–241
longipectoralis 226–241
Etmopterus ***sculptus*** 379–384
Etsaputu 497, 499
relictum 497–502
Gymnorhamphichthys ***britskii*** 400–406
Gymnotus ***tiquie*** 77–81
Hemigrammus ***diagonicus*** 211–215
Himantolophus ***stewarti*** 490–493, 495–496
Micracanthicus 216–225
vandragti 216–225
Neocaristius 385–399
Paracaristius
aquilus 385–399
nemorosus 385–399
nudarcus 385–399
Paulasquama 46–52, 222
callis 46–52, 216, 221, 224
Photonectes 201–210
barnetti 201–210
coffea 201–210
corynodes 201–210
Poecilia ***hondurensis*** 418–422
Scorpiodoras ***liophysus*** 121–140
†***Sharfia*** 64–71
mirabilis 64–71
Siagonodon ***acutirostris*** 58–62
Soromonichthys 216–225
stearleyi 216–225
Steindachnerina ***seriata*** 524–528

NEW AND ESTABLISHED NAMES

Abudefduf
lorenzi 329
natalensis 329
Acacia 424
Acanthicus 216, 220, 224
hystrix 224
Acanthocybium solandri 156
Acanthodoras 121
calderonensis 121
cataphractus 121
Acanthopterygii 590, 592, 594–595
Acanthuridae 317, 582, 585
Acari 569
Acipenser oxyrinchus 157
Acris crepitans 305
Adinia 247
Adrianichthyidae 559
Aeshna sp. 145
Agamyxis 121
Agkistrodon contortrix 556
Albula 593
Alligator
mississippiensis 609
sinensis 608–609
Alosa
aestivalis 334
pseudoharengus 334
sapidissima 334, 336
Alveolina dainelli 66
Amblydoras 121–122, 137
affinis 138
bolivariensis 122, 138
Ambystoma 15, 313
annulatum 5
jeffersonianum 9–17, 530–538
macrodactylum 161
maculatum 9–17
opacum 9–17
tiginum 305
tigrinum 530, 533
Amiiformes 613
Amniota 474
Amphibia 251–263, 513–522
Amphibolurus muricatus 42
Amphiliidae 477–489
Amphilophus citrinellus 317–318
Amphisbaenidae 615
Amphisbaena 615
Anablepidae 243
Anadoras 121–122, 137
grypus 138
weddellii 138
Anaxyrus fowleri 369
Ancistrinae 50
Ancistrini 216–225, 497–502
Ancistrus 46, 50, 222, 224, 497, 500
macrophthalmus 219–220
pirareta 221, 224, 498, 501
Andersonia 478–481
Andropogon sp. 30
Anguilliformes 592
Anolis 41–42
carolinensis 40
cristatellus 41–42
sagrei 38–45
Anomalopus
leuckarti 115
leuckartii 117
Anotophysi 177
Anthiinae 357–364
Anthias
anthias 358–359

© 2011 by the American Society of Ichthyologists and Herpetologists DOI: 10.1643/OT-11-165

nicholsi 358–359
noeli 358–359
Antipodocottus 172
Anura 251–263
Aparasphenodon 258
Amphiuma 313
Apicomplexa 470
Apistogramma 4
Aplocheilichthyinae 243–244
Aplocheilidae 243–244
Aplocheilioidei 243–244
Aprolepis barbarae 99
Araneae 569
Arenophryne 427
rotunda 426
Argentiformes 593
Artemisia filifolia 30
Arvicolinae 431
Aspidoscelis 35
sexlineatus 34
Astroblepus 507
Astrochelys radiata 278, 283
Astrodoradinae 121–122, 137
Astrodoras 121–122, 137
asterifrons 138
heckeli 122
sp. 138
Astyanax metae 215
Athene noctua 548
Atherinidae 247
Atheriniformes 249, 590, 593
Atherinomorpha 242–250, 592, 595
Atherinops affinis 312
Atherinopsidae 247, 249, 308–314, 626
Atherinopsis californiensis 312
Atractosteus
spatula 612
tropicus 613
Atractus 407–417
elaps 410–412, 415
Austrofundulus 313
Autanadoras 121–122
milesi 121–122

Banksia prionotes 424
Baryancistrus
beggini 220
demantoides 220
Batrachochytrium dendrobatidis 302
Batrachoidiformes 592
Belonoglanis 478–481
Beryciformes 592
Blanus 615
Blarina brevicauda 431, 433, 438
Bokermannohyla 258
Bolbometopon muricatum 317–318
Bolitoglossa 457
adspersa 457
altamazonica 457
colonnea 458, 460
nicefori 457–459
paraensis 457–462
rostrata 457–460
rufescens 458–460
subpalmata 460
Bothidae 329
Bothus swio 329
Bouteloua
dactyloides 30
gracilis 30
Brachyhypopomus sp. 79
Brachyraphis 372–378
parismina 372–378
Brycon 177
meeki 581
melanopterus 215
Bufo 304
americanus 444
boreas 141
Bullisichthys 360
caribbeus 358, 360
Buteo buteo 548

Calyptotis scutirostrum 115, 117
Carapidae 335
Carapus bermudensis 584
Carcharhinus 539
altimus 539
plumbeus 539
Carcharodon 181
Caristiidae 385–399
Caristius 385–386
groenlandicus 385
japonicus 385
macropus 393
maderensis 393
sp. 386, 393
sp. cf. *maderensis* 393, 395
Carludovica rotundifolia 352
Carphophiinae 413
Caryospora 470
Casineria 474
Castilla elastica 352
Casuarina equisetifolia 39
Catostomus
commersonii 170
macrocheilus 169–170
Centropristis 357–364
ocyurus 358–359
philadelphica 358–359
striata 358–359
Cepolidae 593
Ceratophrys ornata 519
Chaenichthyidae 581
Chaenopsidae 587
Chaetostoma 46, 50, 216, 222, 224, 497, 501, 503–512
anomalum 221, 222, 224
cf. *nudirostris* 46
lineopunctatum 503–512
loborhynchos 506, 509
pearsei 499, 501
Chanidae 179
Chanos 177–178, 598
Chapinophis 414
Characidae 211–215
Characiformes 177, 179, 211–215, 523–529
Characoidei 177

†*Charitosomus* 178
Chascanopsetta kenyaensis 329
Chaunus arenarum 304
Cheilidoperca 357–364
Cheilinus undulatus 315–318
Chelidae 75
Chelidoperca hirundinac 358–359
Chelodina
expansa 73, 75
longicollis 73, 75
rugosa 313
Chelonia mydas 471
Chiloglanis 482
Chilopoda 569
Chlorurus microrhinos 317
Choerodon azurio 317
Chromis
randalli 329
rapanui 329
Chrysemys picta 472
Chysipterus 329
rapanui 329
Cichla sp. 338
Cichlasoma 4, 317
Cichlidae 317, 593
Cladophora 145
glomerata 143, 145
Clevelandia 95
Clupeiformes 593, 595
Clupeomorpha 592, 595
Cnemidophorus 34
Cocconeis 145
Codoma 82–83, 86, 89–91
ornata 83–84, 86–87, 89–90
Coelophrys bradburyae 603
Coleoptera 569
Collembola 569
Colostethus 427
inguinalis 352
Colpichthys hubbsi 93, 100
Coluber constrictor 32
Colubroidea 407–417
Congoglanis 477–489
alula 477–489
inga 477–489
sagitta 477–489
Coniophanes 407–408, 410–414
fissidens 410–412, 415
Cophomantini 258
Cordylancistrus 50, 224, 501
platycephalus 216
torbesensis 216
Corvus sp. 548
Corymbophanes 51, 501
Corythomantis 258
greeningi 258
Cottidae 172
Cottus carolinae 5–8
Cranoglanis bouderius 509
Craugastor augusti 369
Creediidae 172
Creteuchiloglanis 226–241
brachypterus 226–241
gongshanensis 226–241
kamengensis 226–241
longipectoralis 226–241
macropterus 226–241
Crinia insignifera 426
Cromeria 178
Crossorhombus howensis 329
Crotalus
horridus 430–442
oreganus
o. abyssus 35
o. helleri 439
o. oreganus 439
viridis
v. helleri 439
v. oreganus 439
v. viridis 32
Cryophis 408, 410–414
hallbergi 407–417
Cryptobranchus
alleganiensis 5–8
Cryptosporidium 470
Ctenothrissiformes 592
Ctenotus
leonhardi 115
leonhardii 117
Culaea inconstans 172
Cuneatus 613
Cunuria crassipes 80
Cuora
serrata 275
trifasciata 275
Curimatidae 523–529
Cyclanthaceae 352
Cyclothone 206
Cyprinella 82–83, 86, 89–91
bocagrande 84, 87
caerulea 84
formosa 84, 87, 89–90
lutrensis 84, 87–90
ornata 86, 89–90
proserpina 84, 87
venusta 84, 88
Cyprinidae 89
Cypriniformes 177, 179, 593
Cyprinodontidae 242–243
Cyprinodontiformes 242–250
Cyprinodontoidea 243
Cyprinodontoidei 242–250
Cyrtopodion kotschyi 546
Czatkobatrachus polonicus 611

Danio 178
rerio 169, 563
Dekeyseria 50
niveata 224
Delturus 501
Denticeps 595
Dentilepisosteus 613
Derhamia 50
Dermaptera 569
Desmognathus ochrophaeus 110
Deuterodon 215
Diadophis 408, 414
punctatus 410–412, 415

Dibranchus 603–604
Dinilysia 615
Diplectrum 357–364
bivittatum 358–359
formosum 358–359
macropoma 358–359
pacificum 358–359
rostrum 358–359
Dipodomys ordii 34
Diponoi 616
Dipsadidae 407–417
Dipsadinae 413–414
Dipsas 407–417
catesbyi 410–412, 415
Diptera 569
Dolichancistrus 50, 216, 224, 501
pediculatus 51
Doradidae 121–140
Doras
armatulus 121
(Astrodoras) 133
calderonensis 121–122, 133, 135
carinatus 138
cataphractus 121
depressus 121, 133
fimbriatus 121
Hancoki 122
heckelii 122, 133
micropoeus 138
punctatus 121
(Rhinodoras) 133
Doumea 477–489
alula 477–489
typica 479
Doumeinae 477–489
Dunkleosteus 470
Dyscophus antongilii 611

Echiodon dawsoni 335
Eigenmannia sp. 79
Eimeria 470
Elaphe quatuorlineata 548
Electrophorus 179
electricus 400
Eleutherodactylus coqui 18–28, 369
Elopiformes 592
Elopomorpha 595
Elops 593
Emblemaria 587
pandionis 587
Emblemariopsis 587
Emydidae 606
Enhydrina schistosa 556
Engyprosopon
mozambiquensis 329
regani 329
Engystomops 513
Enneacanthus 467–468
chaetodon 467
gloriosus 467
obesus 463–469
†*Eosladenia* 69
caucasica 64

Epictia 53, 59
albipuncta 61
australis 61
goudotti 61
magnamaculata 61
munoai 61
nasalis 58–59, 61
tenella 61
Epinephelidae 357
Epinephelides 360
armatus 360
Epinephelus 361
armatus 358
Epithemia 145
Eremiascincus 114
richardsoni 115
richardsonii 117
Eretmochelys imbricata 73, 75
†*Erfoudichthys* 179
rosae 179
Ericaceae 598
Eulamprus amplus 115, 117
Etheostoma 319–326
baileyi 319–326
caeruleum 319
coosae 319–326
(Nanostoma) 319
(Ulocentra) 319
zonale 319–326
zonistium 319–326
Etmopteridae 379–384
Etmopterus 379–384
brachyurus 379–384
cf. *molleri* 382
lucifer 379–384
molleri 379–380, 382–383
sculptus 379–384
sp. 379
Etsaputu 497, 499
relictum 497–502
Euchilichthys 482
Euchiloglanis 227–228
feae feae 227, 237
gonghanensis 232
hodgarti 235
kamengensis 234, 236–237
macrotrema 235
Eucyclogobius 95
Eukaryota 473
Eupemphix 513
fuscomaculatus 513–522
nattereri 518–519
Euphorbiaceae 80
Eurycea 109
Euteleostei 179
Evermannia 95
Exastilithoxus 50, 216–217, 224
Exostoma 227–228

Falco
eleonorae 546–548
tinnunculus 548
Farlowella 497
Formicidae 569
Franciscodoras marmoratus 121
Fundulidae 242–250
Funduloidea 243

Fundulus 242, 247
bermudae 249
capensis 242–250
(Fontinus) 247
heteroclitus 242–250, 313
luciae 249
nisorius 242
(Zygonectes) 247

Gadiformes 592
Gambusia affinis 298
Gasterosteus aculeatus 324
Gastropoda 569
Geocrinia 426
alba 427
vitellina 427
Gerobatrachus 475
Gila
cypha 346
elegans 346
robusta 346
Gillichthys 93, 95–100
detrusus 93–102
mirabilis 93–102
seta 96–100
Ginglymostoma cirratum 539–544
Glandulocauda 181
Glaphyromorphus 113–120
douglasi 114–115, 117
fuscicaudis 115, 117
isolepis 114
gracilipes 113–120
Glaridoglanis 227–228
Glaucomys sabrinus 431, 433
Glyphoglossus molossus 9
Glyptosternia 228
Glyptosternina 227
Glyptosternon 227–228
Gobiesociformes 592
Gobiidae 93–102, 247, 361
Gobionellidae 99
Gobius
detrusus 99
townsendi 99
Gobiusculus flavescens 324
Gophonema 145
Gonorynchiformes 177–179, 590, 592
Gonorynchus 178, 598
Goodeidae 243
Gopherus
agassizii 78, 281–283, 607
polyphemus 285
Graptemys
gibbonsi 606
pearlensis 606
Gymnogobius 95
Gymnorhamphichthys 400–406
bogardusi 400, 404
britskii 400–406
cf. *hypostomus* 401, 404
hypostomus 400–406
petiti 404
rondoni 400–401, 404
rosamariae 404–405
sp. 401, 404

Gymnotidae 77–81
Gymnotiformes 77–81, 177, 179, 593
Gymnotocinclus 50
anosteos 50
Gymnotoidei
Gymnotus 77–81
anguillaris 77–81
carapo 77–81
chaviro 77–81
chimarrao 77–81
coatesi 79–80
coropinae 77–81
curupira 77–81
esmeraldas 77
henni 77–81
inaequilabiatus 77–81
javari 79–80
jonasi 79–80
maculosus 77–81
melanopleura 79–80
onca 79–80
panamensis 79–80
pantanal 77–81
pantherinus 77–81
pedanopterus 79–80
sp. 2 78
stenoleucus 79–80
tiquie 77–81
varzea 77–81

Haematoloechus sp. 304
†*Hakeliosomus* 178
Halieutichthys 604
Halieutopsis 603–604
margaretae 603
Heleioporus 424, 427
albopunctatus 423–429
eyrei 426
Heliconia 351–352
Heliconeacae 352
Helicoprion 470
Hemanthias
leptus 358–359
peruanus 358–359
vivanus 358–359
Hemerocoetinae 172
Hemiancistrus
guahiborum 220
maracaiboensis 498, 501
subviridis 220
Hemidactylium scutatum 624
Hemidactylus turcicus 546
Hemiergis 113–120
decresiensis 113–120
d. continentis 118
d. davisi 118
d. decresiensis 118
d. talbingoensis 118
gracilipes 119
initialis 114, 117–119
i. brookeri 115
i. initialis 115
millewae 114–115, 117–119
peronii 114–115, 117–119

quadrilineata 114
quadrilineatum 113–120
talbingoensis 119
Hemigrammus
diagonicus 211–215
erythrozonus 214
gracilis 214
marginatus 215
mimus 214
silimoni 211–215
Hemisus 612
Hepatozoon 301–307
catesbianae 303
clamatae 303
clamitae 301–307
Heterandria formosa 242, 244, 246, 248–249
Heterodon 408, 415
nasicus nasicus 32
platirhinos 443–456
platyrhinos 410–412, 415
Himantolophus 490–496
appelii 490–496
sp. 493–494
stewarti 490–493, 495–496
Himantolophidae 490–496
Hiodon 177, 179
Hisonotus 50
Holbrookia 35
maculata 34
Holocentrus 599
Holostei 590, 612–613
Homopus signatus signatus 278–284
Hopliancistrus 50
Hoplias malabaricus 327
Hyla regilla 535
Hylidae 251–263
Hylinae 257
Hylini 258
Hymenoptera 569
Hypancistrus 216–217
contradens 217, 220
lunaorum 217, 220
Hyphessobrycon 215
Hypnus 95
Hypodoras 121–122, 137
forficulatus 138
Hypoplectrus 357–364
chlorurus 358–359
huntii 358–359
macullochi 358–359
nigricans 358–359
puella 358–359
unicolor 358–359
Hypopomidae 79
Hypopomus artedi 405
Hypopygus
lepturus 79
sp. 79
Hypostominae 216, 222, 497–502
Hypostomini 497
Hypostomus
cf. *plecostomus* 498, 501
squalinus 220
Hypsiboas 258
albopunctatus 258
crepitans 258
raniceps 258
Hypsiglena 407–417
affinis 410–412, 415
chlorophaea 410–412, 415
jani 410–412, 415
ochrorhyncha 410–412, 415
platyrhinos 410–412, 415
slevini 410–412, 415
sp. *Cochise* 410–412, 415
torquata 410–412, 415

Ichthyodectidae 592
Ichthyostega 616
Ictalurus punctatus 156, 161, 509
Iguana iguana 614, 623
Imantodes
cenchoa 408, 410–412, 415
inornatus 408, 410–412, 414–415
Iracema 400
caiana 405
Isopondyli 590
Isoptera 569
Isospondyli 593, 595
Isospondylii 593
Istiompax indica 157
Istiophoridae 153
Istiophorus platyperus 157
Itapotihyla 258
langsdorffi 258

Kajikia audax 153–160
Kalyptodoras bahiensis 121
Kinosternon
arizonense 606
flavescens 606
Kneria 593
Kneriidae 179

Labeo 482
Labeotropheus fuelleborni 324
Labridae 315–318, 361
Lacerta trilineata 546
Lacertidae 545–552
Lampridiformes 335
Lampris guttatus 335
Lampropeltis triangulum 32
Lanius senator 548
Lankesterella 302
Lapemis hardwickii 556
Larus michahellis 546
Lasiancistrus 50, 497, 501
schomburgkii 497
Latimeria 613, 616
Leguminosae 79
Lepidogobius 95
seta 95, 98
Lepidoperca aurantia 358, 360–361
Lepidoptera 569
Lepidosauria 614
Lepisosteidae 613
Lepisosteiformes 612–613
Lepisosteus
oculatus 613
osseus 612

Lepomis 467
macrochirus 467
Leporacanthicus 216–217, 224
galaxias 220
Leposoma guianense 566
Lepospondyli 474
Leptoancistrus 50, 216, 224, 501
Leptodeira 407–417
nigrofasicata 408, 410–412, 414–415
septentrionalis 408, 410–412, 415
Leptodeirini 407–417
Leptolucania 247
Leptotyphlopidae 53–63
Leptotyphlops 53
albifrons 53
borrichianus 53
brasiliensis 53–54, 56–58
cupinensis 53
dulcis 53, 61
goudotii phenops 61
humilis 53
h. boettgeri 53
melanotermus 53
nasalis 53
septemstriatus 53, 57
tesselatus 53
Lepus californicus 472
Leuresthes tenuis 308–314
Lissamphibia 474
Lithobates 305
areolatus 285–295
capito 285–295
catesbeianus 162, 166, 302, 369
clamitans 304, 369
(Nenirana) 285–295
palustris 285–295
pipiens 301–307
sevosus 285–295
sphenocephalus 290
sylvatica 305
sylvaticus 366, 369
Lithodoras 121
Lithogenes 501
Lithoxus 50, 216–217, 224
Liuperus biligonigerus 518
Lobodon carcinophagus 581
Lobotes surinamensis 156
Lonchocarpus urucu 79
Lophiidae 64–71
Lophiiformes 64–71, 490–496, 592
Lophiodes 64–71
caulinaris 64
monodi 64
spilurus 64
Lophiohylini 258
Lophiomus 69
setigerus 64
Lophius
americanus 64, 623
†brachysomus 64–71
Loricaria †plecostomus 64–65
Loricariidae 46–52, 216–225, 497–512
Lucania 247
goodei 324
Luvarus imperialis 585
Lymantria dispar 433
Lyomeri 593
Lythrurus 82–83
ardens 85, 87–88, 90
lirus 85, 87
rosepinnis 85, 88
snelsoni 85, 88

Makaira nigricans 157
Mallotus villosus 312
Masticophis flagellum 32
Mazarunia 50
Megalancistrus 216–217
Meiolania 75
platyceps 72–76
Meiolaniidae 76
Melaleuca 424
Melanostomiidae 201–210
Menidia 248
beryllina 248–249
menidia 248–249, 298
Merodoras 121–122, 137
nheco 138
Metynnis 332
Micracanthicus vandragti 216–225
Micropterus
dolomieu 161–168
salmoides 7
Microtus 438
pennsylvanicus 431, 433, 436
pinetorum 431, 433
Miripinnati 593
Misgurnus
anguillicaudatus 296–300
fossilis 298
Moenkhausia 215
costae 215
lepidura 215
Mollienesia
petenensis 421
sphenops 243
s. altissima 421
s. macrura 421
s. vantynei 421
Moraceae 352
Mormyriformes 592–593
Morone saxatilis 334
Muraenidae 361
Musa 351–352
Musaceae 352
Myersglanis 227–228
Myobatrachidae 423
Myodes 438
gapperi 430–442
Myxine mcmillanae 329
Myxinidae 329
Myxocyprinus 179

Naso
brachycentron 317
fageni 317–318
tuberosus 317
unicornis 318
vlamingii 318

Natrix natrix 548
Neblinichthys 50
Necturus maculosus 531
Negaprion brevirostris 539
Nemodoras 138
Neobatrachus 426
  pelabatoides 426
Neocaristius 385–399
  heemstrai 385–399
Nerodia rhombifer 553–558
Notacanthiformes 592
Notechis scutatus 556
Notemigonus 86
Nothopsis rugosus 408, 413
Notropis 83, 89
  baileyi 85
  emiliae 91
  longirostris 83, 85, 87–90
  maculatus 82–84, 86–91
  scabriceps 89
  simus 89
  stilbius 85, 87, 90
Noturus gyrinus 626

Obaichthyidae 613
Obaichthys 613
Odontophrynus americanus 517
Oedipus estheri 457
Ogcocephalidae 603
Oncorhynchus
  mykiss 6, 161
  nerka 170
  tshawytscha 170
Oneirodes bradburyae 603
Onychomys leucogaster 34
Oophaga pumilio 351–356
Opiliones 569
Opisthognathidae 593
Opsopoeodus 82–92
  emiliae 82–92
Opuntia 472
  sp. 30
Oreoglanis 226–228
Orthoptera 569
Oryzias latipes 559–565
Ostariophysi 177, 179–180, 590, 592, 594–595
Osteichthyes 593
Osteoglossiformes 590, 592–594
Osteoglossomorpha 592, 595
Otophysi 177, 590
Owstoniidae 593
Oxydoras eigenmanni 121

Pachyrhachis 615
Palaeobatrachus 611
Paludicola 517
  albifrons 518
  signifera 516
Panaque 50, 216–217, 497
  albomaculatus 497
  gnomus 497
  (Panaque) 216
Paracanthopterygii 592, 595
Paracaristius 385–399
  aquilus 385–399
  heemstrai 386, 392
  maderensis 385–399
  nemorosus 385–399
  nudarcus 385–399
Parachiloglanis 227–228
Paralabrax 357–364
  auroguttatus 358–359
  clathratus 358–359
  humeralis 358–359
  maculatofasciatus 358–360
  nebulifer 358–360
Paralichthys 247
  sp. 247
Paraphractura 478
Pareuchiloglanis 226–241
  anteanalis 239
  feae 226–227, 229–230, 237–240
  f. feae 226–227
  f. myzostoma 226
  gongshanensis 226–241
  gracilicaudata 226–230, 239
  kamengensis 226–241
  longicauda 226, 239
  macropterus 226–241
  macrotrema 226, 231, 240
  myzostoma 226, 240
  poilanei 226–241
  rhabdurus 240
  robusta 240
  sinensis 226
  songdaensis 240
  tianquanensis 240
Paulasquama 46–52, 222
  callis 46–52, 216, 221, 224
Peckoltia furcata 498, 501
Pederpes 474
Pelobate fuscus 426
Pelophylax 304
Perca flavescens 7
Perciformes 315–318, 592
Percomorpha 385–399
Percophidae 172
Percopsiformes 592
Perognathus
  flavescens 34–35
  flavus 34
Peromyscus 430–442
  leucopus 430, 433, 435, 437–438
  maniculatus 34, 430, 433, 437–438
Pterodoras 121
Pholidophorus 470
Photonectes 201–210
  achirus 202, 204, 206–208
  albipennis 202–203, 206, 208
  braueri 202, 208
  barnetti 201–210
  caerulescens 202, 204, 206–208
  coffea 201–210
  corynodes 201–210
  (Dolichostomias) 202
  dinema 202, 208
  fimbria 206
  gracilis 202, 208
  leucospilus 202
  margarita 201–202, 208

(Melanonectes) 202
mirabilis 202–207, 209
munificus 202
parvimanus 202, 206, 209
(Photonectes) 201–210
phyllopogon 202–204, 206, 208–209
(Trachinostomias) 202, 206
Phractolaemus 179, 593
Phractura 478–481
Phrynosoma 413
Physalaemus 513, 517–518, 521
albifrons 513, 517–519
biligonigerus 513–522
fuscomaculatus 513, 518
marmoratus 513, 518–519
santafecinus 513, 518–519
Physopyxis 121–122, 137
ananas 138
cristata 138
lyra 138
Pimephales 83, 86, 89
notatus 84, 86–91
promelas 84, 86–91
tenellus 85–91
vigilax 85–91
Pinus rigida 431, 435, 438
Pituophis catenifer 32
Plancterus 247
Plasmodium falciparum 583
Platyberyx 385–386
heemstrai 385
opalescens 385
Platycephalidae 361
Plethodon 103, 109–111, 623
albagula 103–112
cinereus 110
glutinosus 103, 109, 110
hoffmani 110
kentucki 110
larselli 110
metcalfi 110
neomexicanus 110
ouachitae 110
sequoyah 109
serratus 110
teyahalee 110
vehiculum 110
yonahlosee 110
Plethodontidae 457–462
Pleurodema 513–522
bibroni 513–522
borellii 514, 516–518, 520
brachyops 517, 520
bufoninum 517, 520
cinereum 517
cordobae 514, 516–518, 520
darwinii 517
diplolister 517, 520
fuscomaculatum 513
guayapae 517, 520
kriegi 513–522
marmoratum 517, 520
nebulosum 517, 520
sp. 516
thaul 517, 520
tucumanum 514, 517–518, 520
Podarcis 546
gaigeae 545–552
milensis 550
Poecilia 418–422
(Allopoecilia) 418
boesemani 419–420
butleri 418–420
catemaconis 419–420
caucana 418–420
caudata 421
chica 419–420
formosa 419–420
gillii 419–421
hondurensis 418–422
koperi 418–420
kykesis 419–421
latipinna 419–421
latipunctata 419–420
mexicana 418–421
marcellinoi 419–421
maylandi 419–420
mechthildae 419–420
(Mollenesia) 418–422
nelsoni 419–420
orri 419–421
petenensis 419–420
reticulata 319, 419–420
rositae 419–420
salvatoris 420–421
sphenops 418–421
sulphuraria 419–420
tenuis 421
teresae 419–420
thermalis 419–420
tropicus 421
vandepolli 419–420
velifera 419–420
vetiprovidentiae 421
vivipara 249, 419–420
Poeciliidae 242–250, 418–422
Poeciliinae 242–250
Poeciliopsis turrabarrensis 375
Poeciloidea 243
Poecilopsetta vaynei 329
Poecilopsettidae 329
Polypterus 613
Pomacentridae 329, 361, 593
Pothos sp. 19
Probolodus 215
Procatopodinae 243–244, 249
Procatopus
schioetzi 249
similis 249
Profundulidae 243
Pronotogrammus martinicensis 358–359
Protacanthopterygii 177, 590, 592, 594–595
Psectrogaster curviventris 527
Pseudacanthicus 216–217
Pseudacris
maculata 365–371
regilla 141

Pseudancistrus 50, 499
anthrax 220
barbatus 51
orinoco 220
tigris 220
Pseudanthias
bicolor 358–359
conspicuus 358–359
elongatus 358–359
squamipinnis 358–359
taeniatus 358–359
Pseudexostoma 226–228
Pseudis 257–258
caraya 258
Pseudochromidae 361
Pseudoleptodeira 407–417
latifasciata 407–408, 410–412, 415
uribei 408
Pseudophryne 313
bibronii 312
Pseudorhombus 247
Pseudoscorpiones 569
Psychrolutidae 172
Pteraclis 285
fasciatus 285
macrops 285
Pterygoplichthyini 497, 499
Pterygoplichthys 497
Ptychocheilus 346
Pungitius pungitius 170
Pygocentrus 181
nattereri 181

Quercus 431
Quietula 95, 247
Quiscalus lugubris 571

Rana 285, 301–307
areolata 289
a. areolata 291
aurora 167
boylii 151–152,161–168
cascadae 161
catesbeiana 162, 530, 535
draytonii 141
erythraea 427
mucosa 161
pipiens 530–531, 533, 535
pretiosa 426
sylvatica 426–427, 530–538, 623
Rena 53
boettgeri 58
humilis 58
unguirostris 61
Rhabdias ranae 304
Rhabdinaea 408, 410–412, 414
fulvivittis 410–412, 415
Rhabdinophanes 414
Rhamphichthyidae 400–406
Rhamphichthys 400
cf. *marmoratus* 405
drepanium 405
hahni 405
Rhinelepini 497
Rhinichthys osculus 161–168
Rhinoderma darwinii 612
Rhinophrynus 612
dorsalis 611
Rineloricaria formosa 224
Rivulidae 243–244

Sacura margaritacea 358–359
Salmoniformes 590, 593
Salvelinus fontinalis 161
Samaridae 329
Samaris
costae 329
desoutterae 329
nielseni 329
Sarcocystis 470
Scaphirhynchus platorynchus 156
Scaridae 317, 473
Scarthyla 258
Sceloporus 35
undulatus 34–35
Schindleria 593
Sciaenops ocellatus 334
Scinax 251–263
acuminatus 251–263
albicans 252, 255–256, 259, 261
alter 251–263
argyreornatus 252, 255–256, 258–259, 261
ariadne 252, 255–256, 258–259, 261
aromothyella 252, 255, 259, 261
auratus 258
berthae 252, 255–256, 258–259, 261
breini 256
canastrensis 251–263
catharinae 251–263
crospedospilus 258
curicica 251–263
cuspidatus 258
duartei 251–263
elaeochrous 256, 258
eurydice 253, 256–257, 260–261
fuscomarginatus 251–263
fuscovarius 251–263
granulates 251–263
hayii 251–263
hiemalis 252, 255–256, 259, 261
littoralis 252, 255–256, 258–259, 261
longilineus 252, 256, 259, 261
nasicus 251–263
obtriangulatus 252, 256, 258–259, 261
perereca 251–263
perpusillus 251, 253, 255–256, 259, 262
rizibilis 253, 255–256, 258–259, 261
rostratus 251, 256
ruber 251–263
similis 251–263
sp. 1 253, 255–256, 262
sp. 2 253, 256, 259, 262
squalirostris 251–263
staufferi 256, 258
trapicheiroi 253, 255–257, 262
uruguayus 251–263
Scincella lateralis 114–117
Scincidae 113–120
Sciuridae 433

Sciurus carolinensis 431–433, 438
Scleroparei 593
Scopelarchoides nicholsi 206
Scorpaeniformes 592
Scorpiodoras 121–140
calderonensis 121–140
cf. *calderonensis* 133
heckelii 121–140
liophysus 121–140
sp. 122
Sebastes carnatus 626
Serpentes 53–63, 614
Serraniculus 357–364
pumilio 358–359
Serranidae 357–364
Serraninae 357
Serranus 357–364
annularis 358–359
atricauda 358–359
baldwini 358–360
cabrilla 358–359
fasciatus 360
hepatus 358–359
notospilus 358–359
phoebe 358–359
psittacinus 358–360
scriba 358–359
subligarius 358–359
tabacarius 358–359
tigrinus 358–359
tortugarum 358–359
†***Sharfia*** 65, 68–69
mirabilis 64–71
Siagonodon 53–63
acutirostris 58–62
borrichianus 58–62
brasiliensis 53–63
cupinensis 60–61
semistriatus 57–62
Sialis sp. 145
Sibon 407–417
nebulatus 410–412, 415
Siluriformes 46–52, 121–140, 177, 179, 216–225, 226–241, 477–489, 497–512, 593
Similiparma 329
Sineoamphisbaena hexatabularis 615
Sisoridae 226–241
Sistrurus
catenatus 29–37, 435, 443–456, 553–558
c. catenatus 33–35
c. edwardsii 29–37, 435
c. tergeminus 34
miliarius 553–558
Skiotocharax 50
meizon 50
Sladenia 64–71
remiger 64
Sorex 431
cinereus 433, 438
Soricidae 433
Soromonichthys stearleyi 216–225
†*Spaniodon hakelensis* 178
Sparidae 361
Spatuloricaria sp. 220
Spectracanthicus 216–217, 499
murinus 50, 499
Spermophilus
spilosoma 34
tridecemlineatus 34
Sphaenorhynchus 258
Sphaerodactylus 566–576
levinsi 566
macrolepis 572
nicholsi 566, 575
parvus 572
townsendi 575
vincenti 566–576
v. ronaldi 566–576
Sphenodon 614
Sphenomorphus 113–114, 116, 118
fasciatus 114
variegatus 114
Sphyrna 539
tiburo 539
Spinax lucifer 379
Squaliformes 379–384
Squamata 113–120
Steindachnerina 523–529
amazonica 524, 527–528
argentea 524, 527–528
atratoensis 524, 527–528
bimaculata 524, 527–528
binotata 524, 527–528
biornata 524, 527–528
brevipinna 523–529
conspersa 524, 527–528
corumbae 523–524
dobula 524, 527–528
elegans 524, 527–528
fasciata 523–529
gracilis 524, 527–528
guentheri 524
guntheri 527–528
hypostoma 524, 527–528
insculpta 524, 527–528
leucisca 524, 527–528
notograptos 524, 527–528
notonota 524, 527–528
planiventris 524, 527–528
pupula 527–528
quasimodoi 524, 527–528
runa 523
seriata 524–528
varii 524, 527–528
Sternopygidae 79
Sternopygus macrurus 79
Stomiidae 201–210
Stomiiformes 201–210
Sturisoma nigrirostrum 497
Stylephorus 338
Subholostei 590
Sylvilagus floridanus 432–433
Symphysanodon haraldi 135
Synentognathi 590

Tamasciurus hudsonicus 431–432
Tamias striatus 431, 433, 438
Tamiasciurus hudsonicus 433, 438

Tantalophis 408, 410–411, 414
discolor 410–412, 415
Taricha
granulosus 141, 161–168
sierra 141
torosa 141
Teleostei 64–71, 93–102, 201–210, 357–364, 385–399, 418–422
Telescopus fallax 548
Temnospondyli 474
Terrapene 275, 472
carolina 270–277, 607
ornata 270–277
o. luteola 283
Tetragonopterinae 211, 213
Tetraodontiformes 473, 584, 588
Tetrapoda 616
Tetrapodomorpha 616
Thamnophis
proximus 553–558
radix 32, 553–558
sirtalis 187–200, 436, 553–558
s. annectens 187
s. pallidus 187
s. parietalis 187–200
s. sirtalis 187–200
Thayeria 215
Thunnus 268
albacares 264–269
obesus 267–268
orientalis 264–269
thynnus 267–268
Thysanoptera 569
Tomeurus gracilis 249
Toxoplasma 470
Trachydoras 121, 138
Trachyglanis 478–481
Triadobatrachus 611
Tricheilostoma 53–63
brasiliensis 54–62
fuliginosum 61
salgueiroi 61
Trichonotidae 172
Trigonophrys rugiceps 519
Trionychidae 606
Triportheus pictus 215
Trogonophidae 615
Trypanosoma 302
Tyto alba 436

Urodela 457–462

Valenciidae 243
Valencioidea 243
Varanus 471

Werthemieria maculata 121

Xenodontinae 413
Xenohyla 258
Xenopus laevis 530–531
Xiphias gladius 155, 157
Xiphophorus 324
milleri 249
Xyrauchen 346

Zamenis situlus 548
Zapus hudsonicus 432–433

SUBJECT INDEX

Copeia 2011, Nos. 1–4

ACTIVITY, *Cottus carolinae* (ontogen. response to predators; foraging) 5–8; *Eleutherodactylus coqui* (calling) 18–28; *Sistrurus catenatus edwardsii* (seas. movement patterns) 29–37; *Anolis sagrei* (displays; communication signals) 38–45; *Rana boylii* (tadpole swim. perform.) 141–152; *Rana boylii* (tadpole response to predators) 161–168; *Lithobates areolatus* (burrow dwell.) 285–295; *Oophaga pumilio* (female territoriality) 351–356; *Crotalus horridus* (forag.) 430–442; *Sistrurus catenatus, Heterodon platirhinos* (movement; spatial dispersion) 443–456; *Gingtymostoma cirratum* (polyandry) 539–544; *Sphaerodactylus vincenti ronaldi* (reprod. seas.; feed.) 566–576.

AGE, *Kajikia audax* (fin-spine annuli; problems) 153–160; *Sphaerodactylus vincenti ronaldi* (skeletochronology) 566–576.

ALARM SIGNALS, *Cryptobranchus alleganiensis* (response of *Cottus carolinae*) 5–8; *Anolis sagrei* (signal rates; agonistic) 38–45.

BEHAVIOR, *Cottus carolinae* (ontogenetic shift, avoid. behav. => forag. behav.) 5–8; *Ambystoma* (*opacum, jeffersonianum, maculatum*) (agonistic; btwn./among spp.) 9–17; *Sistrurus catenatus edwardsii* (hibernacula & forag. site choice) 29–37; *Anolis sagrei* (displays; communication signals) 38–45; *Rana boylii* (tadpole response, diff. water velocities) 141–152; *Rana boylii* (tadpole response, native/non-native predators) 161–168; *Homopus s. signatus* (female reprod. response to rainfall) 278–284; *Lithobates areolatus* (use of crawfish burrows) 285–295; *Leuresthes tenuis* (delayed hatch.) 308–314; *Etheostoma* (*baileyi, coosae, zonale, zonistium*) (nuptial color., spectral diffs.) 319–326; *Oophaga pumilio* (female/female aggression) 351–356; *Heleioporus albopunctatus* (survival; reprod.) 423–429; *Crotalus horridus* (forag.; prey select.) 430–442; *Sistrurus catenatus, Heterodon platirhinos* (movement; spatial dispersion; site fidelity) 443–456; *Gingtymostoma cirratum* (polyandry) 539–544; *Podarcis gaigeae* (reprod.; cannibalism; "island syndrome") 545–552; *Sphaerodactylus vincenti ronaldi* (reprod.; feed.; popn. struct.) 566–576.

BIODIVERSITY, *Gymnotus* (*tiquie* n. sp.) (syntopic spp.) 77–81; *Plethodon albagula* (cryptic lineages) 103–112; *Scorpiodoras* spp. (resurrection, *calderonensis*) 121–140; *Thunnus* (*albacares, orientalis*) (nucl. DNA popn. study) 264–269; *Terrapene* (*carolina, ornata*) (hybrid., introgress.) 270–277; *Etmopterus* (prob. undescribed spp.) 379–384; Caristiidae (partial fam. revision) 385–399; *Poecilia* (*Mollienesia*) (key to Centr. Amer. spp.) 418–422; *Congoglanis* (n. gen., *alula* n. comb., *inga* n. sp., *sagitta* n. sp.) (expansion of Doumeinae) 477–489; *Himantolophus* (*appelii* gr.) (distinct morphs) 490–496; *Etsaputu* (*relictum* n. gen., n. sp.) (Ancistrini tribe) 497–502; *Chaetostoma* (species richness; vicariant event) 503–512; *Steindachnerina* (2nd most species-rich genus w/in Curimatidae) 523–529.

BIOGEOGRAPHY, *Sistrurus catenatus edwardsii* (seas. movement) 29–37; *Paulasquama* (*callis* n. gen., n. sp.) (falls as barrier to derived taxa invasion) 46–52; †*Sharfia* (*mirabilis* n. gen., n. sp.) (Eocene, Monte Bolca; Tethys Sea) 64–71; Meiolaniid (fossil terres. turtle; Miocene; New Zealand vicariant; Oligocene drowning, Moa's Ark) 72–76; *Gillichthys* (*detrusus, mirabilis, seta*) (Colorado R. delta, Baja Calif., no. Gulf of Calif.) 93–102; *Plethodon albagula* (popns., Edwards Plateau, TX; historic) 103–112; *Hemiergis* spp. (distrib. of lineages) 113–120; *Scorpiodoras* spp. (suggest. historic barrier to dispersal) 121–140; *Thamnophis sirtalis* (color var. w/in popns.) 187–200; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (centr. Pacific O.) 201–210; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (W Guiana Shield) 216–225; *Creteuchiloglanis* (Yunnan distrib. compared w/ *Pareuchiloglanis*) 226–241; *Etmopterus* (*sculptus* n. sp.) (endemic, so. Afr. upper cont. slope) 379–384; *Chaetostoma lineopunctatum* (Fitzcarrald Arch; vicariant event) 503–512.

BLOOD, *Lithobates pipiens* (leukocyte profiles, pesticides, parasites) 301–307; *Pleurodema* (*borelli, tucumanum, bibroni, cordobae*) (erythrocytes, table) 513–522.

CALLS, *Eleutherodactylus coqui* (clinal var.; native/non-native) 18–28; *Pseudacris maculata* (compar. index values, reliability) 365–371.

CANNIBALISM, *Ambystoma* (*opacum, jeffersonianum, maculatum*) (agonistic behav. to conspecifics) 9–17; *Podarcis gaigeae* (rel. to gigantism & decr. island size) 545–552.

CHEMORECEPTION, *Cottus carolinae* (response to amphib. alarm cues; ontogen. shift) 5–8; *Rana boylii* (tadpole detect. native/non-native predators) 161–168.

COLORATION, *Paulasquama* (*callis* n. gen., n. sp.) (in life; color photo.) 46–52; *Tricheilostoma brasiliensis* (in life; b&w photo.), *Siagonodon* (*acutirostris* n. sp.) (in preserv.; b&w photo.) 53–63; *Gymnotus* (*tiquie* n. sp.) (in alcohol; b&w photo.) 77–81; *Gillichthys* (*detrusus, mirabilis, seta*) (in life; compar.; b&w photo.) 93–102; *Scorpiodoras* (*calderonensis, heckelii, liophysus* n. sp.) (in life & in alcohol; b&w/color photos.) 121–140; *Thamnophis sirtalis* (erythrism, extreme var. w/in popns.; color photos.) 187–200; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (in preserv.; b&w photos.) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (in alcohol; b&w photos.) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (in life & in alcohol; b&w photos.) 216–225; *Creteuchiloglanis* (*brachypterus* n. sp., *longipectoralis* n. sp.) (in life; b&w photos.) 226–241; *Fundulus capensis* (=*heteroclitus*) (in alcohol; orig. descrip.) 242–250; *Etheostoma* (*baileyi, coosae, zonale, zonistium*) (nuptial color.; spectral characs., graphs) 319–326; *Etmopterus* (*sculptus* n. sp.) (in life; b&w photo., fig.) 379–384; *Gymnorhamphichthys* (*britskii* n. sp.) (in alcohol; b&w photos.; disting. from congeners) 400–406; *Poecilia* (*hondurensis* n. sp.) (in life & in alcohol; b&w photos.) 418–422; *Congoglanis* (n. gen.) (*alula* n. sp.) (in life), (*inga* n. sp., *sagitta* n. sp.) (in alcohol), (b&w photos.) 477–489; *Etsaputu* (*relictum* n. gen., n. sp.) (in life & in alcohol) (characteristic pattern) 497–502; *Steindachnerina* (*seriata* n. sp.) (in life & in alcohol; b&w photo.; pattern similar to *S. fasciata*) 523–529.

COMMUNICATION, *Eleutherodactylus coqui* (calling, clinal var.; native/non-native) 18–28; *Anolis sagrei* (displays, signal rates) 38–45; *Pseudacris maculata* (compar. index values, reliability) 365–371.

COMMUNITY, *Ambystoma* (*opacum, jeffersonianum, maculatum*) (agonistic behav. w/in larv. guild struct.) 9–17; *Gymnotus* (*tiquie* n. sp.) (syntopic gymnotiforms) 77–81; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (other loricariids collected) 216–225.

COMPETITION, *Ambystoma* (*opacum, jeffersonianum, maculatum*) (larv. intra-guild predation; agonistic behav.) 9–17.

CONSERVATION, *Sistrurus catenatus edwardsii* (popn. study) 29–37; *Rana boylii* (tadpole survival wrt dam-flow) 141–152; *Heleioporus albopunctatus* (survival, modif. agric. areas) 423–429; *Sistrurus catenatus, Heterodon platirhinos* (movement/spatial dispersion & effect of roads) 443–456; *Enneacanthus obesus* (distrib. wrt environ. variables) 463–469.

DEEP-SEA, *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (Pacific O.; mesopelagic) 201–210; *Etmopterus* (*sculptus* n. sp.) (so. Afr., upper cont. slope, 450–900 m depth) 379–384; *Neocaristius heemstrai* (n. gen., n. comb.) (So. Hemisphere; mesopelagic, 780 m depth), *Paracaristius* (4 spp.) (worldwide, tropical/subtropical; mesopelagic) 385–399; *Himantolophus* (*stewarti* n. sp.) (so. oceans; meso-/bathypelagic, most 400–1413 m depth) 490–496.

DENTITION, *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (var. length; fig.) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (photo., SEM of jaws) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (jaw morph., poss. diet) 216–225.

© 2011 by the American Society of Ichthyologists and Herpetologists DOI: 10.1643/OT-11-164

DEVELOPMENT, *Leuresthes tenuis* (delayed hatch.; heterokairy) 308–314; *Cheilinus undulatus* (forehead hump morph.) 315–318; *Rana sylvatica, Ambystoma jeffersonianum* (larval corticosterone levels; stress response) 530–538; *Oryzias latipes* (egg production, incubation & hatching) 559–565; *Sphaerodactylus vincenti ronaldi* (growth rate; seasonality) 566–576.

DIET, *Sistrurus catenatus edwardsii* (rodents, lizards) 29–37; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (poss. insectivores) 216–225; *Homopus s. signatus* (herbivory) 278–284; *Crotalus horridus* (prey composition, table; var. btwn. popns.) 430–442; *Etsaputu* (*relictum* n. gen., n. sp.) (prob. epilithic periphyton) 497–502; *Sphaerodactylus vincenti ronaldi* (inverts.; generalist; seas. var.) 566–576.

DISEASE, *Lithobates pipiens* (pollution, lowered immunity, blood, parasites, leukocyte profiles) 301–307.

DISPERSAL, *Sistrurus catenatus edwardsii* (seas. movement patterns) 29–37; Meiolaniid (fossil terres. turtle; Miocene vicar. event; Oligocene drowning, Moa's Ark) 72–76; *Gymnorhamphichthys* (*britskii* n. sp.) (barrier elimination) 400–406; *Sistrurus catenatus, Heterodon platirhinos* (movement & effects of roads) 443–456.

DISTRIBUTION, *Eleutherodactylus coqui* (native, Puerto Rico; introd., Hawaii) 18–28; *Sistrurus catenatus edwardsii* (SW U.S.A. & NE Mex.) 29–37; *Anolis sagrei* (SE U.S.A., Bahamas, Caribb.) 38–45; *Paulasquama* (*callis* n. gen., n. sp.) (type loc., Mazaruni R., Pakaraima Mtns., Guyana; map) 46–52; *Tricheilostoma brasiliensis* (Braz. Cerrado; map), *Siagonodon* (*acutirostris* n. sp.) (Tocantins, Braz. Cerrado; map) 53–63; †*Sharfia* (*mirabilis* n. gen., n. sp.) (Eocene, Monte Bolca) 64–71; Meiolaniid fossil turtle (N.Z.; fw/terres.) 72–76; *Gymnotus* (*tiquie* n. sp.) (trib., Rio Tiquie, Amaz.; map) 77–81; *Gillichthys detrusus* (endemic, Colorado R. Delta; map) 93–102; *Plethodon albagula* (centr. TX to Int. Highlands; map) 103–112; *Hemiergis* (spp. & sspp.) (so. Austr.; map) 113–120; *Scorpiodoras* (Purus Arch barrier), (*calderonensis, heckelii, liophysus* n. sp.) (upper Amaz. Basin, Braz., Venez.; map) 121–140; *Rana boylii* (Angelo Reserve, no. & centr. CA) 141–152; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (Pacific O.; map) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (lower Amaz. floodplain; map) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.) (endemic, lower Ventuari R.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (endemic, Soromoni Creek), (upper Orinoco R., Venez.; map) 216–225; *Creteuchiloglanis* (*brachypterus* n. sp.) (Irrawaddy R., Yunnan), (*kamengensis* n. comb.) (Brahmaputra R. basin, India & China), (*longipectoralis* n. sp.) (upper Mekong R., Yunnan), (*macropterus* n. comb.) (upper Salween R. & upper Irrawaddy R., Yunnan), (*gongshanensis* n. comb.) (upper Salween R., Yunnan) 226–241; *Fundulus capensis* (=*heteroclitus*) (provenance unknown; not So. Afr.) 242–250; *Thunnus* (*albacares, orientalis*) (W Pacific O.) 264–269; *Terrapene* (*carolina, ornata*) (map, current U.S.A. ranges; known hybrid. sites) 270–277; *Misgurnus anguillicaudatus* (ident.; invasive) 296–300; *Brachyrhaphis parismina* (east. lowlands, Costa Rica; map) 372–378; *Etmopterus* (*sculptus* n. sp.) (so. Afr. endem.; upper cont. slope) 379–384; *Neocaristius heemstrai* (n. gen., n. comb.) (So. Hemisphere), *Paracaristius* (*maderensis, nudacrus* n. sp.) (all oceans), (*nemorosus* n. sp., *aquilus* n. sp.) (E Trop. Atl.), (map) 385–399; *Gymnorhamphichthys* (*britskii* n. sp.) (Rio de La Plata, Parana–Paraguay syst. tribs.; map) 400–406; *Poecilia* (*Mollienesia*) (*hondurensis* n. sp. (Caribb. drain. basins, Honduras) 418–422; *Enneacanthus obesus* (disjunct.; Atl. coastal, S to MS) 463–469; *Congoglanis* (n. gen.) (*alula* n. comb.: thru-out Congo R. basin) (*inga* n. sp.: endemic, l. Congo R. near Inga Rapids), (*sagitta* n. sp.: Lualaba R. basin, SE Congo R. system), (map) 477–489; *Himantolophus* (*stewarti* n. sp., *appelii*) (sympatric, so. oceans; map) 490–496; *Etsaputu* (*relictum* n. gen., n. sp.) (relict; restricted, Marañon R., no. Peru; map) 497–502; *Chaetostoma lineopunctatum* (compar. disjunct popns., E Peru) 503–512; *Pleurodema biligonigerus* (amended type locality, Arg.) 513–522; *Steindachnerina* (*seriata* n. sp.) (Rio Tapajós basin, Braz. Amaz.; map) 523–529; *Podarcis gaigeae* (endemic, Skyros Arch., Greece; map) 545–552.

ECOLOGY, *Cottus carolinae* (ontogen. shift in behav.) 5–8; *Ambystoma* (*opacum, jeffersonianum, maculatum*) (larv. guild agonistic behav.; predation, cannibalism) 9–17; *Eleutherodactylus coqui* (calling, clinal var.) 18–28; *Sistrurus catenatus edwardsii* (spatial ecol.; seas. movement; hibernacula; prey; habitat) 29–37; *Anolis sagrei* (communication signals; behav.) 38–45; *Paulasquama* (*callis* n. gen., n. sp.) (habitat; distrib.; life hist.; sex. dimorph.; parasites; eggs) 46–52; *Tricheilostoma brasiliensis, Siagonodon* (*acutirostris* n. sp.) (distribs.) 53–63; †*Sharfia* (*mirabilis* n. gen., n. sp.) (biogeogr.; paleont.; osteol.; habitat) 64–71; Meiolaniid fossil turtle (Miocene; dispersal; distrib.; habitat; biogeogr.) 72–76; *Gymnotus* (*tiquie* n. sp.) (distrib.; gymnotiform community) 77–81; *Gillichthys detrusus* (distrib.; spawn.; habitat) 93–102; *Plethodon albagula* (distrib.; sex. dimorph.) 103–112; *Hemiergis* spp. (distribs.) 113–120; *Scorpiodoras* (*calderonensis, heckelii, liophysus* n. sp.) (distribs.; biogeogr.) 121–140; *Rana boylii* (tadpole survival, swim. perform., growth wrt water velocity) 141–152; *Rana boylii* (tadpole response to predators) 161–168; *Thamnophis sirtalis* (color morphs w/in & btwn. popns.; sex. dimorph.) 187–200; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (distrib.) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (distrib.) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (distrib.; diet; endemism) 216–225; *Creteuchiloglanis* (*gongshanensis* n. comb., *kamengensis* n. comb., *macropterus* n. comb., *brachypterus* n. sp., *longipectoralis* n. sp.) (distrib.) 226–241; *Fundulus capensis* (osteol.; So. Afr. distrib. questioned) 242–250; *Thunnus* (*albacares, orientalis*) (genetic divers.; popn. size) 264–269; *Terrapene* (*carolina, ornata*) (distrib.; hydrib.) 270–277; *Homopus s. signatus* (herbivory; female reprod. response to rainfall; egg production) 278–284; *Lithobates areolatus* (habitat, morphol.) 285–295; *Misgurnus anguillicaudatus* (invasive; distrib.; sex. dimorph) 296–300; *Lithobates pipiens* (lowered immunity; blood parasites, pesticides & leukocyte profiles) 301–307; *Leuresthes tenuis* (delayed hatch.; develop.; heterokairy) 308–314; *Cheilinus undulatus* (forehead hump morph. & develop.; not sex. charac.) 315–318; *Etheostoma* (*baileyi, coosae, zonale, zonistium*) (nuptial color., spectral characs.) 319–326; *Oophaga pumilio* (activity; behav.; female–female interactions, oviposition sites) 351–356; Serranidae (47 spp.) (evol. sex. patterns) 357–364; *Pseudacris maculata* (call rates; popn. ests.) 365–371; *Brachyrhaphis parismina* (life history var. among popns.) 372–378; *Etmopterus* (*sculptus* n. sp.) (distrib.; habitat; endemic) 379–384; *Neocaristius* (n. gen.), *Paracaristius* (distrib.; prob. diet) 385–399; *Gymnorhamphichthys* (*britskii* n. sp.) (distrib.; habitat) 400–406; *Poecilia* (*hondurensis* n. sp.) (distrib.) 418–422; *Heleioporus albopunctatus* (survival, agric. areas; sex ratios; popn. size) 423–429; *Crotalus horridus* (forag. behav. & diet var. btwn. popns.) 430–442; *Sistrurus catenatus, Heterodon platirhinos* (movement/spatial dispersal) 443–456; *Bolitoglossa paraensis* (reprod.; microhabitat use; sex. dimorph.) 457–462; *Enneacanthus obesus* (environ. variables; conservation; distrib.) 463–469; *Congoglanis* (n. gen.) (*alula* n. comb., *inga* n. sp., *sagitta* n. sp.) (distribs.) 477–489; *Himantolophus* (*stewarti* n. sp.) (distrib.; habitat) 490–496; *Etsaputu* (*relictum* n. gen., n. sp.) (distrib.; habitat; diet; relict) 497–502; *Chaetostoma lineopunctatum* (distrib.; vicariance; haplotypes) 503–512; *Steindachnerina* (*seriata* n. sp.) (distrib.; habitat) 523–529; *Rana sylvatica, Ambystoma jeffersonianum* (larv. develop.; corticosterone levels; stress response) 530–538; *Ginglymostoma cirratum* (reprod., polyandry, litter size) 539–544; *Podarcis gaigeae* ("island syndrome;" distrib.; habitat, gigantism; endemics; reprod. biol.) 545–552; *Sistrurus* (*catenatus, miliarius*), *Nerodia rhombifer, Thamnophis* (*proximus, radix, sirtalis*) (sex ratio; fecundity; neonate size & sex. dimorph.) 553–558; *Oryzias latipes* (reprod. biol.; egg production, incubation & hatching; husbandry) 559–565; *Sphaerodactylus vincenti ronaldi* (habitat; diet; reprod. biol.; growth rate; popn. size; prob. predators; caudal autotomy) 566–576.

EGGS, *Paulasquama* (*callis* n. gen., n. sp.) (size) 46–52; *Homopus s. signatus* (effects of drought; female reprod. effort) 278–284; *Leuresthes tenuis* (delayed hatch.; heterokairy) 308–314; *Bolitoglossa paraensis* (clutch size wrt female size; graph) 457–462; *Podarcis gaigeae* (clutch size & volume; egg size; "island syndrome") 545–552; *Oryzias latipes* (reprod. output; incubation, competency; husbandry) 559–565; *Sphaerodactylus vincenti ronaldi* (size; incubation time) 566–576.

ENDANGERED SPECIES, *Sistrurus catenatus edwardsii* (agric. expansion & urban.) 29–37; *Gillichthys detrusus* (restricted habitat) 93–

102; *Rana boylii* (tadpoles & dam-discharge) 141–152; *Rana boylii* (intro. predator; decline) 161–168; *Enneacanthus obesus* (disjunct. distrib.; environ. variables) 463–469.

ENDEMISM/ENDEMICS, Meiolaniid fossil turtle (N.Z. Cenozoic; 1st fw/terres. turtle) 72–76; *Gillichthys detrusus* (no. Colorado R. Delta, Mex.) 93–102; *Hemigrammus* (*diagonicus* n. sp.) (Balaro & Piranha lakes, Braz.) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.) (lower Ventuari R., Venez.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (Soromoni Creek, Venez.) 216–225; *Etmopterus* (*sculptus* n. sp.) (so. Afr., upper cont. slope, depth 450–900 m) 379–384; *Congoglanis* (*inga* n. gen., n. sp.) (type loc.; Inga Falls, l. Congo R.) 477–489; *Etsaputu* (*relictum* n. gen., n. sp.) (relict, Amaz. tribs., Andes Mtns., no. Peru) 497–502; *Steindachnerina* (*seriata* n. sp.) (headwaters, Rio Jamanxim, Braz.), (*fasciata, brevipinna*) (expanded ranges) 523–529; *Podarcis gaigeae* (Skyros Arch., Greece) 545–552.

EVOLUTION, *Hemiergis* spp. (hypoth., limb reduction) 113–120; Serranidae (47 spp.) (sex. patterns; phylogen. anal.) 357–364.

FECUNDITY, *Bolitoglossa paraensis* (clutch size) 457–462; *Podarcis gaigeae* (clutch vol. & size) 545–552; *Sistrurus* (*catenatus, miliarius*), *Nerodia rhombifer, Thamnophis* (*proximus, radix, sirtalis*) (litter size) 553–558; *Oryzias latipes* (reprod. output; husbandry) 559–565; *Sphaerodactylus vincenti ronaldi* (clutch size) 566–576.

FEEDING, *Cottus carolinae, Rana boylii* (tadpoles; decr. w/ incr. water velocity) 141–152; *Homopus s. signatus* (herbivory) 278–284; Dipsadidae (goo-eater group) 407–417; *Podarcis gaigeae* (cannibalism) 545–552; *Sphaerodactylus vincenti ronaldi* (inverts.; seas. var.; table) 566–576.

FOSSIL, †*Sharfia* (*mirabilis* n. gen., n. sp.) (Eocene; Monte Bolca) 64–71; Meiolaniid (N.Z. terres. turtle; Cenozoic) 72–76.

GENETICS, *Opsopoeodus emiliae* (mtDNA & nucl. DNA; homoplasy) 82–92; *Gillichthys* (*detrusus, mirabilis*) (cyt.*b* & nucl. DNA) 93–102; *Hemiergis* spp. (Bayesian methods; mt-rRNA; phylogen. rels.) 113–120; *Scinax* (*catharinae* clade, *ruber* clade) (karyology; cytogenetics; morphometric anal.) 251–263; *Thunnus* (*albacares, orientalis*) (nucl. DNA, microsat. anal.; popn. size) 264–269; *Terrapene* (*carolina, ornata*) (microsat., mtDNA; hybrid.) 270–277; Serranidae (47 spp.) (literature data, sex. patterns) 357–364; Dipsadidae (mtDNA, nucl. DNA, phylogen. anal.) 407–417; *Chaetostoma lineopunctatum* (cyt.*b*; haplotypes, disjunct popns.) 503–512; *Ginglymostoma cirratum* (microsat. DNA; polyandry; litter paternity) 539–544.

GEOGRAPHIC LOCALITIES,

Argentina, *Scinax* (*catherinae* clade, *ruber* clade) 251–263; *Gymnorhamphichthys* (*britskii* n. sp.) 400–406.

Atlantic Ocean, *Neocaristius heemstrai* (n. gen., n. comb.), *Paracaristius* (*nudarcus* n. sp., *maderensis, aquilus* n. sp., *nemorosus* n. sp.) 385–399; (S) *Himantolophus* (*stewarti* n. sp.) 490–496.

Australia, (NSW, Vict., SA, WA) *Hemiergis* (*decresiensis, millewae, peronii, quadrilineatum, initialis*) 113–120; (WA) *Heleioporus albopunctatus* 423–429.

Bahamas, *Anolis sagrei* 38–45.

Brazil, (Cerrado) *Tricheilostoma brasiliensis, Siagonodon* (*acutirostris* n. sp.) 53–63; (Amaz.) *Gymnotus* (*tiquie* n. sp.) 77–81; (Amaz.) *Scorpiodoras* (*calderonensis, heckelii, liophysus* n. sp.) 121–140; (Para) *Hemigrammus* (*diagonicus* n. sp.) 211–215; *Scinax* (*catherinae* clade, *ruber* clade) 251–263; *Gymnorhamphichthys* (*britskii* n. sp.) 400–406; (Para) *Bolitoglossa paraensis* 457–462; (Para) *Steindachnerina* (*seriata* n. sp., *fasciata, brevipinna*) 523–529.

California, *Rana boylii* 141–152, 161–168; *Leuresthes tenuis* 308–314.

Canada, (Man.) *Thamnophis sirtalis* 187–200; (Que.) *Lithobates pipiens* 301–307; (Ont.) *Sistrurus catenatus, Heterodon platirhinos* 443–456.

Central African Republic, *Congoglanis* (*alula* n. comb., *inga* n. sp.) 477–489.

China, (Yunnan) *Creteuchiloglanis* (*brachypterus* n. sp., *longipectoralis* n. sp., *macropterus* n. comb., *gongshanensis* n. comb., *kamengensis* n. comb.) 226–241.

Colorado, *Sistrurus catenatus edwardsii* 29–37; *Pseudacris maculata* 365–371.

Congo, Dem. Rep., *Congoglanis* (*alula* n. comb., *inga* n. sp.) 477–489.

Costa Rica, *Oophaga pumilio* 351–356; *Brachyrhaphis parismina* 372–378.

Florida, *Anolis sagrei* 38–45; *Ginglymostoma cirratum* 539–544; *Sistrurus miliarius* 553–558.

Greece, *Podarcis gaigeae* 545–552.

Guyana, *Paulasquama* (*callis* n. gen., n. sp.) 46–52.

Hawaii, *Eleutherodactylus coqui* 18–28.

Honduras, *Poecilia* (*Mollienesia*) (*hondurensis* n. sp.) 418–422.

Idaho, *Misgurnus anguillicaudatus* 296–300.

India, *Creteuchiloglanis* (*kamengensis* n. comb.) 226–241.

Indian Ocean, *Neocaristius heemstrai* (n. gen., n. comb.), *Paracaristius* (*nudarcus* n. sp., *maderensis*) 385–399; (S) *Himantolophus* (*stewarti* n. sp., *appelii*) 490–496.

Italy, (Monte Bolca) †*Sharfia* (*mirabilis* n. gen., n. sp.) 64–71.

Kentucky, *Etheostoma* (*zonale, baileyi*) 319–326.

Martinique, *Sphaerodactylus vincenti ronaldi* 566–576.

Maryland, *Enneacanthus obesus* 463–469.

Mexico, *Gillichthys* (*detrusus, mirabilis, seta*) 93–102.

Michigan, *Thamnophis sirtalis* 187–200.

Missouri, *Cottus carolinae, Cryptobranchus alleganiensis* 5–8; *Sistrurus catenatus, Nerodia rhombifer, Thamnophis* (*proximus, radix, sirtalis*) 553–558.

Mozambique, *Etmopterus* (*sculptus* n. sp.) 379–384.

Namibia, *Etmopterus* (*sculptus* n. sp.) 379–384.

New Jersey, *Crotalus horridus* 430–442.

New Zealand, (St. Bathans Fauna) Meiolaniid fossil turtle 72–76.

Oregon, *Rana boylii, Micropterus dolomieu* 161–168.

Pacific Ocean, (SW) *Kajikia audax* 153–160; (centr. N, W) *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) 201–210; (W) *Thunnus* (*albacares, orientalis*) 264–269; (W) *Cheilinus undulatus* 315–318; *Neocaristius heemstrai* (n. gen. n. comb.), *Paracaristius* (*nudarcus* n. sp., *maderensis*) 385–399; (S) *Himantolophus* (*stewarti* n. sp., *appelii*) 490–496.

Pennsylvania, *Ambystoma* (*opacum, jeffersonianum, maculatum*) 9–17; *Crotalus horridus* 430–442.

Peru, *Etsaputu* (*relictum* n. gen., n. sp.) 497–502; *Chaetostoma lineopunctatum* 503–512.

Puerto Rico, *Eleutherodactylus coqui* 18–28.

South Africa, *Homopus s. signatus* 278–284; *Etmopterus* (*sculptus* n. sp.) 379–384.

Tennessee, *Etheostoma* (*coosae, zonistium*) 319–326.

Texas, *Plethodon albagula* 103–112; *Terrapene* (*carolina, ornata*) 270–277.

Uruguay, *Scinax* (*catherinae* clade, *ruber* clade) 251–263.

Venezula, *Scorpiodoras heckelii* 121–140; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) 216–225.

Zambia, *Congoglanis* (*sagitta* n. sp.) 477–489.

GIGANTISM, *Podarcis gaigeae* ("island syndrome") 545–552.

GROWTH, *Rana boylii* (tadpoles & effect water velocity) 141–152; *Kajikia audax* (sclerochronology) 153–160; *Lithobates* (4 spp.) (ontogeny, allometry) 285–295; *Leuresthes tenuis* (delayed hatching; heterokairy) 308–314; *Cheilinus undulatus* (forehead hump develop.) 315–318; *Podarcis gaigeae* ("island syndrome;" gigantism) 545–552; *Oryzias latipes* (somatic; von Bertalanffy growth func.) 559–565; *Sphaerodactylus vincenti ronaldi* (rate; seasonality; food intake) 566–576.

HABITAT, *Ambystoma* (*opacum, jeffersonianum, maculatum*) (ephemeral ponds) 9–17; *Sistrurus catenatus edwardsii* (hibernacula & forag. sites) 29–37; *Paulasquama* (*callis* n. gen., n. sp.) (clear-water; cobble & gravel) 46–52; *Tricheilostoma brasiliensis* (open cerrado) 53–63; †*Sharfia* (*mirabilis* n. gen., n. sp.) (inner shelf, Tethys Sea) 64–71; Meiolaniidae fossil turtle (Miocene; fw/terres., N.Z.) 72–76; *Gymnotus* (*tiquie* n. sp.) (terra firma forest streams) 77–81; *Gillichthys detrusus* (lowtide; silty, delta), *mirabilis* (estuaries; muddy), *seta* (high intertidal; rocky/sandy) 93–102; *Rana boylii* (tadpoles in dam./undam. rivers) 141–152; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (mesopelagic) 201–210; *Micracanthicus* (*vandragti* n. gen., n. sp.) (rocky reef), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (riffles, cobble) 216–225; *Lithobates areolatus* (crawfish burrow usage; photo.) 285–295; *Misgurnus anguillicaudatus* (effect on sex ratio) 296–300; *Oophaga pumilio* (vegetation parameters & female territoriality) 351–356; *Brachyrhaphis parismina* (life hist. var. btwn. clades)

372–378; *Etmopterus* (*sculptus* n. sp.) (upper cont. slope, 450–900 m depth) 379–384; *Neocaristius heemstrai* (n. gen., n. comb.), *Paracaristius* (*maderensis, nudacrus* n. sp., *nemorosus* n. sp., *aquilus* n. sp.) (mesopelagic; most >800 m depth) 385–399; *Gymnorhamphichthys* (*britskii* n. sp.) (sm. streams to lg. rivers) 400–406; *Heleioporus albopunctatus* (survival in agric. areas) 423–429; *Crotalus horridus* (var. & effect on forag. behav.) 430–442; *Sistrurus catenatus, Heterodon platirhinos* (effect of roads on movement/dispersal) 443–456; *Bolitoglossa paraensis* (microhabitat use) 457–462; *Enneacanthus obesus* (environ. variables; agric. & urban. impacts) 463–469; *Congoglanis* (n. gen.) (*alula* n. comb., *sagitta* n. sp., *inga* n. sp.) (rheophilic) 477–489; *Himantolophus* (*stewarti* n. sp.) (meso-/bathypelagic, most 400–1413 m depth) 490–496; *Etsaputu* (*relictum* n. gen., n. sp.) (high-gradient, cobble substrate) 497–502; *Steindachnerina* (*seriata* n. sp.) (flooded area, dense aq. veget.) 523–529; *Podarcis gaigeae* (island endemics; "island syndrome") 545–552; *Sphaerodactylus vincenti ronaldi* (xerophytic) 566–576.

HERMAPHRODITISM, *Cheilinus undulatus* (protogyny; male forehead hump develop.) 315–318; Serranidae (47 spp.) (evol. sex patterns; table) 357–364.

HIBERNATION, *Sistrurus catenatus edwardsii* (hibernacula site) 29–37.

HISTORICAL PERSPECTIVES, Joseph S. Nelson 169–176; Dannie Alan Hensley 327–331; John E. Olney 332–341; Robert Rush Miller 342–347; James Chase Tyler, II, 577–589; "Phyletic studies of teleostean fishes" 590–598; Margaret G. Bradbury 599–605.

HOME RANGE, *Sistrurus catenatus edwardsii* (spatial ecol.; seas. movement) 29–37; *Oophaga pumilio* (female territoriality) 351–356.

HUSBANDRY, *Rana sylvatica, Ambystoma jeffersonianum* (larv. hormone levels; rearing conditions) 530–538; *Oryzias latipes* (reprod. biol.; somatic growth) 559–565.

HYBRIDS/HYBRIDIZATION, *Terrapene* (*carolina, ornata*) (molec. anal.; introgress.; color photos.) 270–277.

IMMUNITY, *Lithobates pipiens* (pesticides, lowered resistance; blood parasites) 301–307.

LARVAE, *Ambystoma* (*opacum, jeffersonianum, maculatum*) (guild; agonistic behav.) 9–17; *Rana boylii* (tadpole swim. perform., growth, survival) 141–152; *Rana boylii* (tadpole response to predators) 161–168; *Leuresthes tenuis* (delayed hatch.; growth, develop.) 308–314; *Rana sylvatica, Ambystoma jeffersonianum* (corticosterone levels, develop./metamoph.; stress response) 530–538; *Oryzias latipes* (incubation period; husbandry) 559–565.

LIFE HISTORY, *Cottus carolinae* (ontogen. shift, response to amphib. alarm cues) 5–8; *Ambystoma* (*opacum, jeffersonianum, maculatum*) (larv. guild; behav.) 9–17; *Sistrurus catenatus edwardsii* (seas. movement; hibernacula & forag. habitat) 29–37; *Anolis sagrei* (breed. seas.; communication/signals) 38–45; *Paulasquama* (*callis* n. gen., n. sp.) (eggs, spawn. period; parasites) 46–52; *Gillichthys detrusus* (prob. spawn./reprod. period) 93–102; *Rana boylii* (effect of water velocity on tadpole survival) 141–152; *Rana boylii* (tadpole & pred. threat) 161–168; *Lithobates areolatus* (habitat) 285–295; *Leuresthes tenuis* (delayed hatch.) 308–314; *Cheilinus undulatus* (develop. forehead hump wrt growth, not sex. charac.) 315–318; *Oophaga pumilio* (female territoriality factors) 351–356; Serranidae (47 spp.) (evol. sex. patterns) 357–364; *Brachyrhaphis parismina* (reprod. var. btwn. ciades) 372–378; *Sistrurus catenatus, Heterodon platirhinos* (compar., movement/spatial dispersal; effect of roads) 443–456; *Bolitoglossa paraensis* (reprod. cycle) 457–462; *Podarcis gaigeae* (reprod. biol.; "island syndrome;" gigantism, cannibalism) 545–552; *Oryzias latipes* (reprod. biol.; husbandry) 559–565; *Sphaerodactylus vincenti ronaldi* (age; growth rate; habitat; reprod. biol.; diet) 566–576.

MERISTICS/MORPHOMETRICS, *Paulasquama* (*callis* n. gen., n. sp.) (table) 46–52; *Tricheilostoma brasiliensis, Siagonodon* (*acutirostris* n. sp.) (table, compar. w/ 3 congeners) 53–63; †*Meiolania platyceps* (table, compar. w/ 3 spp.) 72–76; *Gymnotus* (*tiquie* n. sp.) (table, compar. w/ sp. gr.) 77–81; *Gillichthys* (*detrusus, mirabilis*) (table) 93–102; *Plethodon albagula* (tables: male vs. female; mtDNA groups) 103–112; *Scorpiodoras* (*calderonensis, heckelii, liophysus* n. sp.) (table) 121–140; *Photonectes* spp. (table) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (table) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (table) 216–225; *Creteuchiloglanis* (*gongshanensis* n. comb., *kamengensis* n. comb., *macropterus* n. comb., *longipectoralis* n. sp., *brachypterus* n. sp.) (table) 226–241; *Lithobates* (*areolatus, palustris, capito, sevosus*) (table) 285–295; *Etmopterus* (*sculptus* n. sp.) (table) 379–384; *Neocaristius heemstrai* (n. gen., n. comb.), *Paracaristius* (*nemorosus* n. sp., *nudacrus* n. sp., *aquilus* n. sp., *maderensis*) (table) 385–399; *Congoglanis* (n. gen.) (*alula* n. comb., *inga* n. sp., *sagitta* n. sp.) (table) 477–489; *Etsaputu* (*relictum* n. gen., n. sp.) (table) 497–502; *Chaetostoma lineopunctatum* (morphol. diverg. btwn. popns.) 503–512; *Steindachnerina* (*seriata* n. sp.) (table) 523–529.

METHODS, *Sistrurus catenatus edwardsii* (radio telemetry; seas. movement) 29–37; *Opsopoeodus emiliae* (genetic anal.; taxon. relations.) 82–92; *Hemiergis* spp. (Bayesian anal., mt-rRNA) 113–120; *Kajikia audax* (fin-spine aging problems) 153–160; *Fundulus capensis* (x-ray, CT scan; sp. validity) 242–250; *Scinax* (*catharinae* clade, *ruber* clade) (karyology; cytogenetics; systematics) 251–263; *Thunnus* (*albacares, orientalis*) (nucl. DNA anal.; Bayesian ests.) 264–269; *Lithobates* (*areolatus, palustris, capito, sevosus*) (compar. morphometrics) 285–295; *Misgurnus anguillicaudatus* (field diagnostic) 296–300; *Etheostoma* (*baileyi, coosae, zonale, zonistium*) (reflec. spectrophotometry) 319–326; *Pseudacris maculata* (call index values: popn. size reliability) 365–371; Dipsadidae (coalescent-based molec. anal.) 407–417; *Rana sylvatica, Ambystoma jeffersonianum* (stress response/hormone levels; larval rearing conditions) 530–538.

MICROSATELLITE DNA, *Thunnus* (*albacares, orientalis*) (genetic divers.; popn. size anal.) 264–269; *Terrapene* (*carolina, ornata*) (molec. anal. hybrid.) 270–277; Dipsadidae (species-tree anal.) 407–417; *Ginglymostoma cirratum* (polyandry; litter paternity) 539–544.

MIGRATION, *Sistrurus catenatus edwardsii* (seas. movement; forag. & hibernacula sites) 29–37.

MITOCHONDRIAL DNA, *Opsopoeodus emiliae* (cyt.*b* & ND2; taxon. relations.) 82–92; *Gillichthys* (*detrusus, mirabilis, seta*) (cyt.*b*; sp. validity) 93–102; *Terrapene* (*carolina, ornata*) (molec. anal. hybrid.) 270–277; Dipsadidae (species-tree anal.) 407–417.

MORPHOLOGY, *Paulasquama* (*callis* n. gen., n. sp.) (diagn.; descrip.; merist./morph.; b&w/color photos.; sex. dimorph.; osteol.) 46–52; *Tricheilostoma brasiliensis* (rediagn.; hemipenis; geog. var.; merist./morph.; sex. dimorph.; b&w photo., fig.), *Siagonodon* (*acutirostris* n. sp.) (diagn.; holotype descrip.; color.; merist./morph.; b&w photo., fig.) 53–63; †*Sharfia* (*mirabilis* n. gen., n. sp.) (diagn.; descrip.; osteol.; color photos.; figs.) 64–71; Meiolaniid (N.Z. fossil turtle) (pelvis & thoracic vert.; descrip.; b&w photo., fig.), (compar. w/ †*Meiolania platyceps*) 72–76; *Gymnotus* (*tiquie* n. sp.) (diagn.; descrip.; b&w photo.; merist./morph.), (compar. w/ sp. gr.) 77–81; *Gillichthys* (*detrusus, mirabilis, seta*) (descrip. diffs.; b&w/color photos.; merist./morph.) 93–102; *Plethodon albagula* (popn. var. compar.; sex. dimorph.) 103–112; *Scorpiodoras* (redescrip.; spp. key), (*calderonensis, heckelii, liophysus* n. sp.) (diagn.; descrip.; osteol.; gas bladder; b&w/color photos.) 121–140; *Thamnophis sirtalis* (erythrism, color morphs; color photos.) 187–200; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (diagn.; descrip.; barbels, gill filaments, tooth lengths, figd.; b&w photos.) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (diagn.; descrip.; b&w photos.; SEM, teeth) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (diagn.; descrip.; b&w photos.; osteol.) 216–225; Glyptosternina (table, diagn. traits), *Creteuchiloglanis* (diagn.; key to spp.; spp. meristics, table; figs., compar. to *Pareuchiloglanis*), (*gongshanensis* n. comb., *kamengensis* n. comb., *macropterus* n. comb.) (diagn.; b&w photos.), (*longipectoralis* n. sp., *brachypterus* n. sp.) (diagn.; descrip.; b&w photos.) 226–241; *Fundulus capensis* (redescrip.; syn. of *F. heteroclitus*; osteol.; b&w photo.; x-ray, CT scan) 242–250; *Homopus s. signatus* (shell vs. body mass; eggshell dimensions) 278–284; *Lithobates* (*areolatus, palustris, capito, sevosus*) (compar.; x-rays; osteol.) 285–295; *Misgurnus anguilli-*

caudatus (sex. dimorph. P-fin; b&w photos.) 296–300; *Cheilinus undulatus* (forehead hump growth; not sex. dimorph.; photos.) 315–318; *Etheostoma* (*baileyi, coosae, zonale, zonistium*) (nuptial color., spectral characs.) 319–326; *Brachyrhaphis parismina* (life hist. var. btwn. popns.; round- vs. narrow-bodied clades) 372–378; *Etmopterus* (*sculptus* n. sp.) (diagn.; descrip.; color.; b&w photo., fig.), (compar. w/ congeners) 379–384; *Neocaristius heemstrai* (n. gen., n. comb.) (diagn.; redescrip.), *Paracaristius maderensis* (rediagn.; redescrip.), (*nemorosus* n. sp., *nudacrus* n. sp., *aquilus* n. sp.) (diagn; descrip.), (b&w photos., figs.; meristics; key to spp.) 385–399; *Gymnorhamphichthys* (*britskii* n. sp.) (diagn.; descrip.; b&w photos.), (compar. w/ *G. hypostomus*) 400–406; *Poecilia* (*Mollienesia*) (*hondurensis* n. sp.) (diagn.; descrip.; b&w photos.; fig. gonopodium), (compar. w/ Centr. Amer. congeners; key to spp.) 418–422; *Congoglanis* (n. gen.; diagn.), (*alula* n. comb.) (diagn.; redescrip.), (*inga* n. sp., *sagitta* n. sp.) (diagn.; descrip.), (morphometric table; b&w photos.; key to spp.) 477–489; *Himantolophus* (*stewarti* n. sp.) (diagn; descrip.; color photo.; illicia & escae figd.), (compar. w/ *H. appelii*; figd.) 490–496; *Etsaputu* (*relictum* n. gen., n. sp.) (diagn.; descrip.; osteol.; morphometrics; b&w photo.) 497–502; *Eupemphix fuscomaculatus, Pleurodema bibroni, Physalaemus biligonigerus* (compar.; photos.: whole body, head, inguinal region, foot) 513–522; *Steindachnerina* (*seriata* n. sp.) (diagn.; descrip.; color.; morphometrics; b&w photo.), (phylogen. relation. w/in genus; cladogram) 523–529; *Oryzias latipes* (somatic growth) 559–565.

MUSEUMS, †*Sharfia mirabilis* (n. gen., n. sp.) (misident. matl.) 64–71; *Fundulus capensis* (re-ident.; subj. syn. of *F. heteroclitus*) 242–250; *Lithobates* (*areolatus, palustris, capito, sevosus*) 285–295.

NOMENCLATURE, *Paulasquama callis* (n. gen., n. sp.) 46–52; *Tricheilostoma brasiliensis* (n. comb.), *Siagonodon* (*acutirostris* n. sp.) 53–63; †*Sharfia mirabilis* (n. gen., n. sp.) 64–71; *Gymnotus* (*tiquie* n. sp.) 77–81; *Opsopoeodus emiliae* (recommend. generic status) 82–92; *Gillichthys detrusus* (resurrec.) 93–102; *Glaphyromorphus gracilipes* (transfer to *Hemiergis*) 113–120; *Scorpiodoras* (*calderonensis*: resurrec.), (*liophysus* n. sp.) 121–140; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) 201–210; *Hemigrammus* (*diagonicus* n. sp.) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) 216–225; *Creteuchiloglanis* (n. gen.), (*gongshanensis* n. comb., *kamengensis* n. comb., *macropterus* n. comb., *longipectoralis* n. sp., *brachypterus* n. sp.) 226–241; *Fundulus capensis* (subj. syn. of *F. heteroclitus*) 242–250; *Etmopterus* (*sculptus* n. sp.) 379–384; *Neocaristius heemstrai* (n. gen., n. comb.), *Paracaristius* (*nemorosus* n. sp., *nudacrus* n. sp., *aquilus* n. sp.) 385–399; *Gymnorhamphichthys* (*britskii* n. sp.) 400–406; *Poecilia* (*Mollienesia*) (*hondurensis* n. sp.) (table, status of congeners) 418–422; *Congoglanis* (n. gen.), (*alula* n. comb.), (*inga* n. sp., *sagitta* n. sp.), (Doumeinae characs. delimited) 477–489; *Himantolophus* (*stewarti* n. sp.) 490–496; *Etsaputu* (*relictum* n. gen., n. sp.) 497–502; *Eupemphix fuscomaculatus* (jr. syn. of *Physalaemus biligonigerus*) 513–522; *Steindachnerina* (*seriata* n. sp.) 523–529.

NONINDIGENOUS SPECIES, *Eleutherodactylus coqui* (Hawaii) 18–28; *Micropterus dolomieu* (OR; predation on *Rana boylii*) 161–168; *Misgurnus anguillicaudatus* (ident.) 296–300.

NUCLEAR DNA, *Opsopoeodus emiliae* (S7 intron; taxon. relations.) 82–92; *Gillichthys* (*detrusus, mirabilis, seta*) (S7 intron; sp. validity) 93–102; *Thunnus* (*albacares, orientalis*) (anal., genetic divers. & popn. size) 264–269; Dipsadidae (molec. syst.) 407–417.

OLFACTION, *Rana boylii* (chem. cues, tadpole recogn. native/non-native predators) 161–168.

ORIENTATION, *Crotalus horridus* (forag./non-forag. postures; color photos./diagrams) 430–442.

OSTEOLOGY, *Paulasquama* (*callis* n. gen., n. sp.) (opercles, fig.; charac. compar.) 46–52; †*Sharfia* (*mirabilis* n. gen., n. sp.) (descrip., full body; color photos.; figs.) 64–71; Meiolaniid (N.Z. fossil terres. turtle) (pelvis; thoracic vert.; b&w photos., figs.) 72–76; *Scorpiodoras heckelii* (descrip.; photos., figs.) 121–140; *Kajikia audax* (sclerochronology, fin-spine; fig., photo.) 153–160; *Hemigrammus* (*diagonicus* n. sp.) (SEM, teeth) 211–215; *Soromonichthys* (*stearleyi* n. gen., n. sp.) (fig., head plates & superficial bones) 216–225; *Creteuchiloglanis* (premax. tooth patches compar. to *Pareuchiloglanis*) 226–241; *Fundulus capensis* (x-ray, CT scan; compar. to *F. heteroclitus*) 242–250; *Lithobates* (*areolatus, palustris, capito, sevosus*) (compar. morphometrics; x-rays, fig.) 285–295; *Congoglanis* (n. gen.; x-ray, caudal vert.; compar. to *Doumea typica*) 477–489; *Etsaputu* (*relictum* n. gen., n. sp.) (coracoid, opercle, branchiostegals figd.; compar. to ancistrines) 497–502.

PALEONTOLOGY, †*Sharfia* (*mirabilis* n. gen., n. sp.) (Eocene; Monte Bolca) 64–71; Meiolaniid (N.Z. fw/terres. fossil turtle; Moa's Ark; Cenozoic) 72–76; *Chaetostoma lineopunctatum* (Miocene; Fitzcarrald Arch; vicariance; speciation) 503–512.

PARASITES, *Paulasquama* (*callis* n. gen., n. sp.) (gut nematodes) 46–52; *Lithobates pipiens* (*Hepatozoon* spp.; leukocyte profiles) 301–307.

PESTICIDES, *Lithobates pipiens* (lowered immunity) 301–307.

PHYLOGENETIC ANALYSIS, *Paulasquama* (*callis* n. gen., n. sp.) (place. in *Ancistrus* clade) 46–52; *Opsopoeodus emiliae* (mtDNA & nucl. DNA; taxon. relations.) 82–92; *Gillichthys* (*detrusus, mirabilis, seta*) (mtDNA & nucl. DNA; morphol.; sp. validity) 93–102; *Hemiergis* spp., *Glaphyromorphus gracilipes* (mt-rRNA, Bayesian anal.) 113–120; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (positions w/in prev. anal.) 216–225; Serranidae (47 spp.) (evol. sex. patterns) 357–364; Dipsadidae (molec. anal.) 407–417; *Steindachnerina* (*seriata* n. sp.) (morphol.; place. w/in genus) 523–529.

PHYLOGENY, †*Sharfia* (*mirabilis* n. gen., n. sp.) (place. w/in lophiids) 64–71; *Lithobates* (4 spp.) (relationships) 285–295; *Congoglanis* (n. gen.; place. w/in Doumeinae) 477–489; *Etsaputu* (*relictum* n. gen., n. sp.) (basal ancistrine loricariid) 497–502.

PHYSIOLOGY, *Lithobates pipiens* (leukocyte profiles, pesticides, parasites) 301–307; *Leuresthes tenuis* (heterokairy) 308–314; *Rana sylvatica, Ambystoma jeffersonianum* (larval corticosterone levels; develop.; stress response) 530–538; *Oryzias latipes* (reprod. biol.; somatic growth) 559–565; *Sphaerodactylus vincenti ronaldi* (growth rate; reprod. biol.) 566–576.

POLLUTION, *Lithobates pipiens* (pesticides, lowered immunity, parasites) 301–307.

POPULATIONS, *Plethodon albagula* (mtDNA distinct grs.; morph. compar.) 103–112; *Thamnophis sirtalis* (extreme color var. w/in & btwn. popns.) 187–200; *Thunnus* (*albacares, orientalis*) (genetic anal., W Pacific popns.) 264–269; *Pseudacris maculata* (popn. size not related to call index values) 365–371; *Brachyrhaphis parismina* (life hist. var. btwn. popns./clade) 372–378; *Heleioporus albopunctatus* (survival in agric. areas; sex ratios; popn. size) 423–429; *Crotalus horridus* (geogr. var., forag. ecol.) 430–442; *Chaetostoma lineopunctatum* (compar. disjunct popns.; speciation, vicariant event) 503–512; *Podarcis gaigeae* ("island syndrome;" reprod. biol.; gigantism) 545–552; *Sphaerodactylus vincenti ronaldi* (density/distrib.) 566–576.

PREDATION, *Ambystoma* (*opacum, jeffersonianum, maculatum*) (larv. guild.; agonistic behav.) 9–17; *Rana boylii* (tadpole response to native/non-native predators) 161–168; *Sphaerodactylus vincenti ronaldi* (prob. preds.; caudal autotomy freq.) 566–576.

RELICT, *Etsaputu* (*relictum* n. gen., n. sp.) (restrict. distrib., Andes, no. Peru) 497–502.

REPRODUCTION, *Gillichthys detrusus* (prob. spawn. period) 93–102; *Homopus s. signatus* (female reprod. response to drought, egg prod.) 278–284; *Etheostoma* (*baileyi, coosae, zonale, zonistium*) (male breed. colors; spectral anal.; visual ecology) 319–326; Serranidae (47 spp.) (evol. sex. patterns; phylogen. anal.) 357–364; *Brachyrhaphis parismina* (life hist. var., narrow- vs. round-bodied clades; reprod. value hypoth.) 372–378; *Poecilia* (*Mollienesia*) (*hondurensis* n. sp.) (gonopodium figd.) 418–422; *Sistrurus catenatus, Heterodon platirhinos* (dispersion wrt hibernacula) 443–456; *Bolitoglossa paraensis* (characs., male & female; clutch size; seasonality) 457–462; *Gingłymostoma cirratum* (polyandry, litter paternity; microsat. DNA) 539–544; *Podarcis gaigeae* ("island syndrome;" clutch size & vol.; egg size) 545–552; *Sistrurus* (*catenatus, miliarius*), *Nerodia rhombifer, Thamnophis* (*proximus, radix, sirtalis*) (litter sex ratio; fecundity) 553–558; *Oryzias latipes* (egg production, incubation, competency;

fecundity) 559–565; *Sphaerodactylus vincenti ronaldi* (seasonality; egg size; incubation period; age at first maturity) 566–576.

SCLEROCHRONOLOGY, *Kajikia audax* (fin-spine annuli) 153–160; *Sphaerodactylus vincenti ronaldi* (growth rate; age at first maturity) 566–576.

SEX RATIO, *Misgurnus anguillicaudatus* (effect of marginal *vs.* preferred habitat) 296–300; *Heleioporus albopunctatus* (fluctuation btwn. yrs.) 423–429; *Sistrurus* (*catenatus, miliarius*), *Nerodia rhombifer, Thamnophis* (*proximus, radix, sirtalis*) (litters) 553–558.

SEXUAL DIMORPHISM, *Paulasquama* (*callis* n. gen., n. sp.) (nuptial male characs.) 46–52; *Tricheilostoma brasiliensis* (scale & size var.) 53–63; *Plethodon albagula* (centr. TX males w/ larger body size) 103–112; *Thamnophis sirtalis* (no. popn. females more red) 187–200; *Misgurnus anguillicaudatus* (P-fin; field diagnostic) 296–300; *Cheilinus undulatus* (forehead hump not sex. charac.; photos., figs.) 315–318; *Poecilia* (*Mollienesia*) (*hondurensis* n. sp.) (b&w photos.: male, female) 418–422; *Bolitoglossa paraensis* (females larger) 457–462; *Sistrurus* (*catenatus, miliarius*), *Nerodia rhombifer, Thamnophis* (*proximus, radix, sirtalis*) (neonate size) 553–558.

SIZE, *Brachyrhaphis parismina* (maturity) 372–378; *Bolitoglossa paraensis* (female body wrt clutch size) 457–462; *Podarcis gaigeae* (islands, gigantism; egg/clutch size) 545–552; *Sistrurus* (*catenatus, miliarius*), *Nerodia rhombifer, Thamnophis* (*proximus, radix, sirtalis*) (litter) 553–558.

SURVIVAL, *Rana boylii* (tadpoles, effect of water velocity) 141–152; *Leuresthes tenuis* (delayed hatch.) 308–314; *Heleioporus albopunctatus* (agric. areas; sex ratios; popn. size) 423–429.

SWIM BLADDER, *Scorpiodoras* (*heckelli, calderonensis, liophysus*) (morphol.; photo.) 121–140.

SYSTEMATICS, *Paulasquama* (n. gen.) (place. w/in *Ancistrus* clade) 46–52; *Tricheilostoma brasiliensis* (n. comb.), (*Leptotyphlops* history) 53–63; †*Sharfia* (n. gen.) (place. w/in lophiids) 64–71; *Gymnotus* (*tiquie* n. sp.) (compar. w/ sp. gr.) 77–81; *Opsopoeodus emiliae* (taxon. relation to other No. Amer. cyprinid genera) 82–92; *Gillichthys detrusus* (sp. validity) 93–102; *Hemiergis* spp., *Glaphyromorphus gracilipes* (phylogen. rels.) 113–120; *Scorpiodoras* (place. w/in subfam. Astrodoradinae) 121–140; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (place. w/in subgen. *Photonectes*) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (place. w/in "*H. gracilis* gr.") 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (place. w/in tribe Ancistrini) 216–225; *Creteuchiloglanis* (n. gen.; compar. to *Pareuchiloglanis*) 226–241; *Fundulus capensis* (subj. syn. of *F. heteroclitus*) 242–250; *Scinax* (*catharinae* clade, *ruber* clade) (cytogen. data) 251–263; *Etmopterus* (*sculptus* n. sp.) (compar. w/ "*E. lucifer*" gr.) 379–384; Caristiidae (partial revision; key to spp.; n. gen., n. spp.) 385–399; *Gymnorhamphichthys* (*britskii* n. sp.) (compar. w/ *G. hypostomus*) 400–406; Dipsadidae (historic review; molec. anal.) 407–417; *Poecilia* (*Mollienesia*) (key to Centr. Amer. congeners) 418–422; Doumeinae (historic review), *Congoglanis* (n. gen.; place. w/in subfam.; key to spp.) 477–489; *Himantolophus* (*stewarti* n. sp.) (place. w/in *appelii* gr.) 490–496; *Etsaputu* (*relictum* n. gen., n. sp.) (basal place. w/in Ancistrini) 497–502; *Eupemphix fuscomaculatus* (jr. syn. of *Physalaemus biligonigerus*) 513–522; *Steindachnerina* (*seriata* n. sp.) (place. w/in genus; cladogram) 523–529.

TAXONOMY, *Paulasquama* (*callis* n. gen., n. sp.) (descrip.) 46–52; *Tricheilostoma brasiliensis* (n. comb.; redescrip.), *Siagonodon* (*acutirostris* n. sp.) (descrip.) 53–63; †*Sharfia* (*mirabilis* n. gen., n. sp.) (descrip.) 64–71; *Gymnotus* (*tiquie* n. sp.) (descrip.) 77–81; *Gillichthys detrusus* (compar. w/ *mirabilis* & *seta*; validity) 93–102; *Scorpiodoras* (gen. redescrip.), (*calderonensis, heckelii*: redescrips.), (*liophysus* n. sp.: descrip.) 121–140; *Thamnophis sirtalis* (color-based ssp. questionable) 187–200; *Photonectes* (*coffea* n. sp., *barnetti* n. sp., *corynodes* n. sp.) (descrip.), (subgen. characs.; table) 201–210; *Hemigrammus* (*diagonicus* n. sp.) (descrip.) 211–215; *Micracanthicus* (*vandragti* n. gen., n. sp.), *Soromonichthys* (*stearleyi* n. gen., n. sp.) (descrip.) 216–225; *Creteuchiloglanis* (*gongshanensis* n. comb., *kamengensis* n. comb., *macropterus* n. comb., *longipectoralis* n. sp., *brachypterus* n. sp.) 226–241; *Fundulus capensis* (redescrip.; subj. syn. of *F. heteroclitus*) 242–250; *Etmopterus* (*sculptus* n. sp.) (descrip.) 379–384; *Neocaristius heemstrai* (n gen., n. comb.), *Paracaristius* (*maderensis, nemorosus* n. sp., *nudacrus* n. sp., *aquilus* n. sp.) (descrips.; key to spp.) 385–399; *Gymnorhamphichthys* (*britskii* n. sp.) (descrip.; compar. w/ *G. hypostomus*) 400–406; *Poecilia* (*Mollienesia*) (*hondurensis* n. sp.) (descrip.), (compar., key to congeners) 418–422; *Congoglanis* (n. gen.), (*alula* n. comb., *inga* n. sp., *sagitta* n. sp.) (descrips.; key to spp.) 477–489; *Himantolophus* (*stewarti* n. sp.) (descrip.; compar. to *appelii* gr.) 490–496; *Etsaputu* (n. gen.) (diagn.), (*relictum* n. sp.) (descrip.) 497–502; *Eupemphix fuscomaculatus* (jr. syn. of *Physalaemus biligonigerus*) 513–522; *Steindachnerina* (*seriata* n. sp.) (descrip.; compar. w/in genus) 523–529.

TEMPERATURE, *Eleutherodactylus coqui* (effect on calls, clinal var.; native/non-native popns.) 18–28.

VARIATION, *Eleutherodactylus coqui* (calls, clinal var.; native/non-native popns.) 18–28; *Tricheilostoma brasiliensis* (geog. var.) 53–63; *Plethodon albagula* (morph. var. btwn. popns.) 103–112; *Scorpiodoras* (*heckelli, calderonensis, liophysus*) (swimbladder; photo.) 121–140; *Kajikia audax* (fin-spine aging; w/in & among structure var.) 153–160; *Thamnophis sirtalis* (color morphs w/in & btwn. popns.) 187–200; *Brachyrhaphis parismina* (reprod. var. w/in & among popns.) 372–378; *Heleioporus albopunctatus* (agric. areas & popn. size) 423–429; *Crotalus horridus* (geogr. var., forag. ecol.) 430–442; *Podarcis gaigeae* (diff. reprod. output btwn. island popns.) 545–552; *Sphaerodactylus vincenti ronaldi* (reprod. seas.; popn. size & distrib.) 566–576.

VISION, *Rana boylii* (tadpole recogn. native/non-native predators) 161–168.

VOLUME CONTENTS

Copeia 2011, Nos. 1–4

Albert, James S. (*see* **Carvalho, Tiago P.**)
Albert, James S. (*see* **Maxime, Emmanuel L.**)
Anderson, Jason S. The Rise of Amphibians: 365 Million Years of Evolution. By R. L. Carroll ... 473
Armbruster, Jonathan W., and Donald C. Taphorn. A New Genus and Species of Weakly Armored Catfish from the Upper Mazaruni River, Guyana (Siluriformes: Loricariidae) ... 46
Armbruster, Jonathan W. (*see* **Lujan, Nathan K.**)
Baker, R. J. (*see* **Salcedo, N. J.**)
Baldo, Diego (*see* **Cardozo, Dario E.**)
Baldo, Diego (*see* **Kolenc, Francisco**)
Beard, Karen H. (*see* **O'Neill, Eric M.**)
Beckstead, Thomas H. (*see* **Mulcahy, Daniel G.**)
Belden, Lisa K. (*see* **Chambers, David L.**)
Belk, Mark C., Emily E. Nance, and Jerald B. Johnson. Life History of *Brachyrhaphis parismina*: Variation Within and Among Populations ... 372
Birindelli, José L. O. (*see* **Sousa, Leandro M.**)
Black, Ron (*see* **Rouse, Jeremy D.**)
Blanton, Rebecca E., Lawrence M. Page, and Bradley M. Ennis. Phylogenetic Relationships of *Opsopoeodus emiliae*, with Comments on the Taxonomic Implications of Discordance among Datasets ... 82
Blaustein, Andrew R. (*see* **Paoletti, David J.**)
Borteiro, Claudio (*see* **Cardozo, Dario E.**)
Borteiro, Claudio (*see* **Kolenc, Francisco**)
Bortoleto, João F. (*see* **Cardozo, Dario E.**)
Brochu, Christopher A. The Chinese Alligator: Ecology, Behavior, Conservation, and Culture. By J. Thorbjarnarson and X. Wang ... 608
Brooks, Ronald J. (*see* **Rouse, Jeremy D.**)
Buchman, Anna B. (*see* **Cureton, James C., II**)
Bushar, Lauretta M. (*see* **Reinert, Howard K.**)
Cailliet, Gregor M. (*see* **Iwamoto, Tomio**)
Cardozo, Dario E., Daniela M. Leme, João F. Bortoleto, Glaucilene F. Catroli, Diego Baldo, Julián Faivovich, Francisco Kolenc, Ana P. Z. Silva, Claudio Borteiro, Célio F. B. Haddad, and Sanae Kasahara. Karyotypic Data on 28 Species of *Scinax* (Amphibia: Anura: Hylidae): Diversity and Informative Variation ... 251
Carnevale, Giorgio (*see* **Pietsch, Theodore W.**)
Carrier, Jeffrey C. (*see* **Heist, Edward J.**)
Carvalho, Tiago P., Cristiane S. Ramos, and James S. Albert. A New Species of *Gymnorhamphichthys* (Gymnotiformes: Rhamphichthyidae) from the Paraná–Paraguay Basin ... 400
Cashner, Frances Miller (*see* **Cashner, Robert Charles**)
Cashner, Robert Charles, Gerald R. Smith, Gifford Hubbs Miller, Frances Miller Cashner, and Barry Chernoff. Robert Rush Miller (1916–2003) ... 342
Casper, Gary S. (*see* **Mooi, Randall D.**)
Catroli, Glaucilene F. (*see* **Cardozo, Dario E.**)
Chakrabarty, Prosanta. Fish Karyotypes: A Checklist. By Ryoichi Arai ... 613
Chakrabarty, Prosanta, and William L. Fink. Piranha 3D. Directed by Alexandre Aja ... 181
Chambers, David L., Jeremy M. Wojdak, Pang Du, and Lisa K. Belden. Corticosterone Level Changes throughout Larval Development in the Amphibians *Rana sylvatica* and *Ambystoma jeffersonianum* Reared under Laboratory, Mesocosm, or Free-living Conditions ... 530
Chernoff, Barry (*see* **Cashner, Robert Charles**)
Ciccotto, Patrick J. Water Chemistry and Landscape Patterns in the Distribution of Banded Sunfish (*Enneacanthus obesus*) in Streams of the Atlantic Coastal Plain in Maryland ... 463
Clack, Jennifer A. How Vertebrates Left the Water. By M. Laurin ... 615
Cohen, Daniel M. (*see* **Iwamoto, Tomio**)
Compagno, Leonard J. V. (*see* **Ebert, David A.**)
Corn, Paul Stephen, Erin Muths, Amanda M. Kissel, and Rick D. Scherer. Breeding Chorus Indices Are Weakly Related to Estimated Abundance of Boreal Chorus Frogs ... 365
Courtenay, Walter R., Jr., Thomas A. Munroe, Richard Winterbottom, Ramon Ruiz-Carus, and William F. Smith-Vaniz. Dannie Alan Hensley (1944–2008) ... 327
Curcio, Felipe F. (*see* **Pinto, Roberta R.**)
Cureton, James C., II, Anna B Buchman, Raelynn Deaton, and William I. Lutterschmidt. Molecular Analysis of Hybridization between the Box Turtles *Terrapene carolina* and *T. ornate* ... 270
David, Solomon. An Empirical Synthetic Pattern Study of Gars (Lepisosteiformes) and Closely Related Species, Based Mostly on Skeletal Anatomy. The Resurrection of Holostei. By Lance Grande ... 612
Davie, Peter S. (*see* **Kopf, R. Keller**)
Davis, Drew R., and Gregory B. Pauly. Morphological Variation among Populations of the Western Slimy Salamander on the Edwards Plateau of Central Texas ... 103
Davis, Matthew P., and Sarah Z. Gibson. The Rise of Fishes: 500 Million Years of Evolution. Second edition. By J. A. Long (ed.) ... 470
Davis, Robert A., and J. Dale Roberts. Survival and Population Size of the Frog *Heleioporus albopunctatus* in a Highly Modified, Agricultural Landscape ... 423
De Vries, Marlee J. (*see* **Ebert, David A.**)
Deaton, Raelynn (*see* **Cureton, James C., II**)
Du, Pang (*see* **Chambers, David L.**)
Ebert, David A., Leonard J. V. Compagno, and Marlee J. De Vries. A New Lanternshark (Squaliformes: Etmopteridae: *Etmopterus*) from Southern Africa ... 379
Ellingson, Ryan A. (*see* **Swift, Camm C.**)
Engbrecht, Nathan J., Susan J. Lannoo, John O. Whitaker, and Michael J. Lannoo. Comparative Morphometrics in Ranid Frogs (Subgenus *Nenirana*): Are Apomorphic Elongation and a Blunt Snout Responses to Small-bore Burrow Dwelling in Crawfish Frogs (*Lithobates areolatus*)? ... 285
Ennis, Bradley M. (*see* **Blanton, Rebecca E.**)
Erisman, Brad E., and Philip A. Hastings. Evolutionary Transitions in the Sexual Patterns of Fishes: Insights from a Phylogenetic Analysis of the Seabasses (Teleostei: Serranidae) ... 357

© 2011 by the American Society of Ichthyologists and Herpetologists DOI: 10.1643/OT-11-151

Esch, Mackenzie (*see* **Reinert, Howard K.**)
Faivovich, Julián (*see* **Cardozo, Dario E.**)
Faivovich, Julián (*see* **Kolenc, Francisco**)
Farrell, Terence M. (*see* **Pilgrim, Melissa A.**)
Fáveri, Sarita B. (*see* **Neckel-Oliveira, Selvino**)
Feller, K. D. (*see* **Gumm, J. M.**)
Ferraris, Carl J., Jr., Richard P. Vari, and Paul H. Skelton. A New Genus of African Loach Catfish (Siluriformes: Amphiliidae) from the Congo River Basin, the Sister-Group to All Other Genera of the Doumeinae, with the Description of Two New Species 477
Ferraro, Daiana P. (*see* **Kolenc, Francisco**)
Findley, Lloyd T. (*see* **Swift, Camm C.**)
Fink, William L. (*see* **Chakrabarty, Prosanta**)
Flessa, Karl W. (*see* **Swift, Camm C.**)
Foufopoulos, Johannes (*see* **Pafilis, Panayiotis**)
Galatti, Ulisses (*see* **Neckel-Oliveira, Selvino**)
Gall, Brian G., and Alicia Mathis. Ontogenetic Shift in Response to Amphibian Alarm Cues by Banded Sculpins (*Cottus carolinae*) 5
Gibson, Sarah Z. (*see* **Davis, Matthew P.**)
Godthelp, Henk (*see* **Worthy, Trevor H.**)
Gumm, J. M., K. D. Feller, and T. C. Mendelson. Spectral Characteristics of Male Nuptial Coloration in Darters (*Etheostoma*) 319
Haddad, Célio F. B. (*see* **Cardozo, Dario E.**)
Hand, Suzanne J. (*see* **Worthy, Trevor H.**)
Hartel, Karsten E. (*see* **Parenti, Lynne R.**)
Hastings, Philip A. (*see* **Erisman, Brad E.**)
Heist, Edward J., Jeffrey C. Carrier, Harold L. Pratt, Jr., and Theo C. Pratt. Exact Enumeration of Sires in the Polyandrous Nurse Shark (*Ginglymostoma cirratum*) 539
Henen, Brian T. (*see* **Loehr, Victor J. T.**)
Hilton, Eric J. Gonorynchiformes and Ostariophysan Relationships: A Comprehensive Review. By Terry Grande, Francisco José Poyato-Ariza, and Rui Diogo 177
Hilton, Eric J., G. David Johnson, Edward D. Houde, and Robert J. Latour. John Edward Olney, Sr. (1947–2010) 332
Hofmeyr, Margaretha D. (*see* **Loehr, Victor J. T.**)
Houde, Edward D. (*see* **Hilton, Eric J.**)
Iwamoto, Tomio, Gregor M. Cailliet, Daniel M. Cohen, Theodore W. Pietsch, Tom Tucker, Ralph J. Larson, and Marlene L. Martin. Margaret G. Bradbury (1927–2010) 599
Jacobs, David K. (*see* **Swift, Camm C.**)
Jiao, Yan (*see* **Leaf, Robert T.**)
Johnson, G. David (*see* **Hilton, Eric J.**)
Johnson, Jerald B. (*see* **Belk, Mark C.**)
Kasahara, Sanae (*see* **Cardozo, Dario E.**)
Kenaley, Christopher P. (*see* **Pietsch, Theodore W.**)
Kenaley, Christopher P. (*see* **Stevenson, Duane E.**)
Kissel, Amanda M. (*see* **Corn, Paul Stephen**)
Klepadlo, Cynthia. Three New Species of the Genus *Photonectes* (Teleostei: Stomiiformes: Stomiidae: Melanostomiinae) from the Pacific Ocean 201
Koetsier, Peter (*see* **Urquhart, Alexander N.**)
Kolenc, Francisco, Diego Baldo, Claudio Borteiro, Federico Marangoni, Daiana P. Ferraro, and Julián Faivovich. The Identity of *Eupemphix fuscomaculatus* Steindachner, 1864 (Amphibia: Anura) 513
Kolenc, Francisco (*see* **Cardozo, Dario E.**)
Kopf, R. Keller, and Peter S. Davie. Fin-Spine Selection and Section Level Influence Potential Age Estimates of Striped Marlin (*Kajikia audax*) 153
Kramer, Jeffrey I. (*see* **Leaf, Robert T.**)
Kupferberg, Sarah J., Amy J. Lind, Vanessa Thill, and Sarah M. Yarnell. Water Velocity Tolerance in Tadpoles of the Foothill Yellow-legged Frog (*Rana boylii*): Swimming Performance, Growth, and Survival 141
Lannoo, Michael J. (*see* **Engbrecht, Nathan J.**)
Lannoo, Susan J. (*see* **Engbrecht, Nathan J.**)
Larson, Ralph J. (*see* **Iwamoto, Tomio**)
Latour, Robert J. (*see* **Hilton, Eric J.**)
Leaf, Robert T., Yan Jiao, Brian R. Murphy, Jeffrey I. Kramer, Karl M. Sorensen, and Victor G. Wooten. Life-History Characteristics of Japanese Medaka *Oryzias latipes* 559
Leclair, Maria Helena (*see* **Leclair, Raymond, Jr.**)
Leclair, Raymond, Jr., and Maria Helena Leclair. Life-History Traits in a Population of the Dwarf Gecko, *Sphaerodactylus vincenti ronaldi*, from a Xerophytic Habitat in Martinique, West Indies 566
Lee, Julian C. The Snakes of Honduras: Systematics, Distribution, and Conservation. By James R. McCranie 609
Leme, Daniela M. (*see* **Cardozo, Dario E.**)
Li, Xu (*see* **Zhou, Wei**)
Lima, Amanda A. (*see* **Neckel-Oliveira, Selvino**)
Lima, Crisalda (*see* **Neckel-Oliveira, Selvino**)
Lima, Flávio C. T. (*see* **Maxime, Emmanuel L.**)
Lind, Amy J. (*see* **Kupferberg, Sarah J.**)
Linsenmair, K. Eduard (*see* **Meuche, Ivonne**)
Liu, Min, and Yvonne Sadovy de Mitcheson. Forehead Morphology of the Humphead Wrasse *Cheilinus undulatus* (Perciformes: Labridae) in Relation to Body Size 315
Loehr, Victor J. T., Brian T. Henen, and Margaretha D. Hofmeyr. Reproductive Responses to Rainfall in the Namaqualand Speckled Tortoise 278
Lujan, Nathan K., and Jonathan W. Armbruster. Two New Genera and Species of Ancistrini (Siluriformes: Loricariidae) from the Western Guiana Shield 216
Lujan, Nathan K., Jonathan W. Armbruster, and Blanca Rengifo. A New Basal Ancistrini Genus and Species from the Andes of Northern Peru (Siluriformes: Loricariidae) 497
Lutterschmidt, William I. (*see* **Cureton, James C., II**)
MacGregor, Gylla A. (*see* **Reinert, Howard K.**)
Mackessy, Stephen P. (*see* **Wastell, Andrew R.**)
Maisano, Jessica A. The Skull and Appendicular Locomotor Apparatus of Lepidosauria. Biology of the Reptilia, Volume 21, Morphology I. By C. Gans, A. S. Gaunt, and K. Adler (eds.) 614
Marangoni, Federico (*see* **Kolenc, Francisco**)
Marcogliese, David J. (*see* **Shutler, Dave**)
Maret, Timothy J. (*see* **Mott, Cy L.**)
Martin, Karen L. (*see* **Moravek, Cassadie L.**)
Martin, Marlene L. (*see* **Iwamoto, Tomio**)
Mathis, Alicia (*see* **Gall, Brian G.**)
Maxime, Emmanuel L., Flávio C. T. Lima, and James S. Albert. A New Species of *Gymnotus* (Gymnotiformes: Gymnotidae) from Rio Tiquié in Northern Brazil 77
May, Peter G. (*see* **Pilgrim, Melissa A.**)
McMahan, Caleb D. A Field Guide to Coastal Fishes from Maine to Texas. By Val Kells and Kent Carpenter 472
Mendelson, T. C. (*see* **Gumm, J. M.**)
Mendonça, Marina B., and Wolmar B. Wosiacki. A New Species of *Hemigrammus* from the Lower Amazon Floodplain (Characiformes: Characidae) 211

Meuche, Ivonne, K. Eduard Linsenmair, and Heike Pröhl. Female Territoriality in the Strawberry Poison Frog (*Oophaga pumilio*) 351
Miller, Gifford Hubbs (*see* **Cashner, Robert Charles**)
Miyamoto, Michael M. (*see* **Qiu, Fan**)
Mooi, Randall D., Jonathan P. Wiens, and Gary S. Casper. Extreme Color Variation within Populations of the Common Gartersnake, *Thamnophis sirtalis*, in Central North America, with Implications for Subspecies Status 187
Moravek, Cassadie L., and Karen L. Martin. Life Goes On: Delayed Hatching, Extended Incubation, and Heterokairy in Development of Embryonic California Grunion, *Leuresthes tenuis* 308
Mott, Cy L., and Timothy J. Maret. Species-Specific Patterns of Agonistic Behavior among Larvae of Three Syntopic Species of Ambystomatid Salamanders 9
Mulcahy, Daniel G., Thomas H. Beckstead, and Jack W. Sites, Jr. Molecular Systematics of the Leptodeirini (Colubroidea: Dipsadidae) Revisited: Species-tree Analyses and Multi-locus Data 407
Munroe, Thomas A. (*see* **Courtenay, Walter R., Jr.**)
Murphy, Brian R. (*see* **Leaf, Robert T.**)
Muths, Erin (*see* **Corn, Paul Stephen**)
Nance, Emily E. (*see* **Belk, Mark C.**)
Neckel-Oliveira, Selvino, João F. M. Sarmento, Ulisses Galatti, Pablo Suárez, Crisalda Lima, Amanda A. Lima, and Sarita B. Fáveri. Reproductive Traits of the Brazilian Salamander *Bolitoglossa paraensis* (Urodela: Plethodontidae) 457
Netto-Ferreira, André L., and Richard P. Vari. New Species of *Steindachnerina* (Characiformes: Curimatidae) from the Rio Tapajós, Brazil, and Review of the Genus in the Rio Tapajós and Rio Xingu Basins 523
Nussear, Kenneth E. Turtles: The Animal Answer Guide. By J. W. Gibbons and J. Greene 607
O'Neill, Eric M., and Karen H. Beard. Clinal Variation in Calls of Native and Introduced Populations of *Eleutherodactylus coqui* 18
Olson, Deanna H. (*see* **Paoletti, David J.**)
Pafilis, Panayiotis, Johannes Foufopoulos, Kostas Sagonas, Anna Runemark, Erik Svensson, and Efstratios D. Valakos. Reproductive Biology of Insular Reptiles: Marine Subsidies Modulate Expression of the "Island Syndrome" 545
Page, Lawrence M. (*see* **Blanton, Rebecca E.**)
Paoletti, David J., Deanna H. Olson, and Andrew R. Blaustein. Responses of Foothill Yellow-legged Frog (*Rana boylii*) Larvae to an Introduced Predator 161
Parenti, Lynne R., and Karsten E. Hartel. Osteology Identifies *Fundulus capensis* Garman, 1895 as a Killifish in the Family Fundulidae (Atherinomorpha: Cyprinodontiformes) 242
Pauly, Gregory B. (*see* **Davis, Drew R.**)
Pietsch, Theodore W. Greenwood, Rosen, Weitzman, and Myers: Circumstances Surrounding the Publication of "Phyletic Studies of Teleostean Fishes, with a Provisional Classification of Living Forms," 1966 590
Pietsch, Theodore W., and Giorgio Carnevale. A New Genus and Species of Anglerfish (Teleostei: Lophiiformes: Lophiidae) from the Eocene of Monte Bolca, Italy 64
Pietsch, Theodore W., and Christopher P. Kenaley. A New Species of Deep-sea Ceratioid Anglerfish, Genus *Himantolophus* (Lophiiformes: Himantolophidae), from Southern Waters of All Three Major Oceans of the World 490
Pietsch, Theodore W. (*see* **Iwamoto, Tomio**)
Pilgrim, Melissa A., Terence M. Farrell, Peter G. May, Marcus R. Vollman, and Richard A. Seigel. Secondary Sex Ratios in Six Snake Species 553
Pinto, Roberta R., and Felipe F. Curcio. On the Generic Identity of *Siagonodon brasiliensis*, with the Description of a New Leptotyphlopid from Central Brazil (Serpentes: Leptotyphlopidae) 53
Poeser, Fred N. A New Species of *Poecilia* from Honduras (Teleostei: Poeciliidae) 418
Pratt, Harold L., Jr. (*see* **Heist, Edward J.**)
Pratt, Theo C. (*see* **Heist, Edward J.**)
Pröhl, Heike (*see* **Meuche, Ivonne**)
Qiu, Fan, and Michael M. Miyamoto. Use of Nuclear DNA Data to Estimate Genetic Diversity and Population Size in Pacific Bluefin and Yellowfin Tuna (*Thunnus orientalis* and *T. albacares*) 264
Ramos, Cristiane S. (*see* **Carvalho, Tiago P.**)
Reeder, Tod W., and Jennifer D. Reichert. Phylogenetic Relationships within the Australian Limb-Reduced Lizard Genus *Hemiergis* (Scincidae: Squamata) as Inferred from the Bayesian Analysis of Mitochondrial rRNA Gene Sequences 113
Reichert, Jennifer D. (*see* **Reeder, Tod W.**)
Reinert, Howard K., Gylla A. MacGregor, Mackenzie Esch, Lauretta M. Bushar, and Robert T. Zappalorti. Foraging Ecology of Timber Rattlesnakes, *Crotalus horridus* 430
Rengifo, Blanca (*see* **Lujan, Nathan K.**)
Roberts, J. Dale (*see* **Davis, Robert A.**)
Rodriguez, D. (*see* **Salcedo, N. J.**)
Rouse, Jeremy D., Robert J. Willson, Ron Black, and Ronald J. Brooks. Movement and Spatial Dispersion of *Sistrurus catenatus* and *Heterodon platirhinos*: Implications for Interactions with Roads 443
Ruiz-Carus, Ramon (*see* **Courtenay, Walter R., Jr.**)
Runemark, Anna (*see* **Pafilis, Panayiotis**)
Sadovy de Mitcheson, Yvonne (*see* **Liu, Min**)
Sagonas, Kostas (*see* **Pafilis, Panayiotis**)
Salcedo, N. J., D. Rodriguez, R. E. Strauss, and R. J. Baker. The Fitzcarrald Arch: A Vicariant Event for *Chaetostoma* (Siluriformes: Loricariidae) Speciation? 503
Sarmento, João F. M. (*see* **Neckel-Oliveira, Selvino**)
Scherer, Rick D. (*see* **Corn, Paul Stephen**)
Scofield, R. Paul (*see* **Worthy, Trevor H.**)
Seigel, Richard A. (*see* **Pilgrim, Melissa A.**)
Shutler, Dave, and David J. Marcogliese. Leukocyte Profiles of Northern Leopard Frogs, *Lithobates pipiens*, Exposed to Pesticides and Hematozoa in Agricultural Wetlands 301
Silva, Ana P. Z. (*see* **Cardozo, Dario E.**)
Simon, Valerie B. Communication Signal Rates Predict Interaction Outcome in the Brown Anole Lizard, *Anolis sagrei* 38
Široký, Pavel. The Biology of the Coccidia (Apicomplexa) of Snakes of the World. A Scholarly Handbook for Identification and Treatment. By Donald W. Duszynski and Steve J. Upton 470
Sites, Jack W., Jr. (*see* **Mulcahy, Daniel G.**)
Skelton, Paul H. (*see* **Ferraris, Carl J., Jr.**)
Smith-Vaniz, William F. (*see* **Courtenay, Walter R., Jr.**)

Smith, David. James Chase Tyler, II 577
Smith, David. Joseph Schieser Nelson 169
Smith, Gerald R. (*see* **Cashner, Robert Charles**)
Sorensen, Karl M. (*see* **Leaf, Robert T.**)
Sousa, Leandro M., and José L. O. Birindelli. Taxonomic Revision of the Genus *Scorpiodoras* (Siluriformes, Doradidae) with Resurrection of *Scorpiodoras calderonensis* and Description of a New Species 121
Stephens, Patrick R. Turtles of the United States and Canada. Second edition. By C. H. Ernst and J. E. Lovich 606
Stevenson, Duane E., and Christopher P. Kenaley. Revision of the Manefish Genus *Paracaristius* (Teleostei: Percomorpha: Caristiidae), with Descriptions of a New Genus and Three New Species 385
Stinner, Jerry N. Life in a Shell: A Physiologist's View of a Turtle. By D. C. Jackson 471
Strauss, R. E. (*see* **Salcedo, N. J.**)
Suárez, Pablo (*see* **Neckel-Oliveira, Selvino**)
Svensson, Erik (*see* **Pafilis, Panayiotis**)
Swift, Camm C., Lloyd T. Findley, Ryan A. Ellingson, Karl W. Flessa, and David K. Jacobs. The Delta Mudsucker, *Gillichthys detrusus*, a Valid Species (Teleostei: Gobiidae) Endemic to the Colorado River Delta, Northernmost Gulf of California, Mexico 93
Taphorn, Donald C. (*see* **Armbruster, Jonathan W.**)
Tennyson, Alan J. D. (*see* **Worthy, Trevor H.**)
Thill, Vanessa (*see* **Kupferberg, Sarah J.**)
Thomson, Alfred W. (*see* **Zhou, Wei**)
Trueb, Linda. Frogs. The Animal Answer Guide. By M. E. Dorcas and J. W. Gibbons 611
Tucker, Tom (*see* **Iwamoto, Tomio**)
Urquhart, Alexander N., and Peter Koetsier. Pectoral Fin Morphology as a Reliable Field Sexing Characteristic in Populations of the Invasive Oriental Weatherfish (*Misgurnus anguillicaudatus*) 296
Valakos, Efstratios D. (*see* **Pafilis, Panayiotis**)
Vari, Richard P. (*see* **Ferraris, Carl J., Jr.**)
Vari, Richard P. (*see* **Netto-Ferreira, André L.**)
Vollman, Marcus R. (*see* **Pilgrim, Melissa A.**)
Wastell, Andrew R., and Stephen P. Mackessy. Spatial Ecology and Factors Influencing Movement Patterns of Desert Massasauga Rattlesnakes (*Sistrurus catenatus edwardsii*) in Southeastern Colorado 29
Whitaker, John O. (*see* **Engbrecht, Nathan J.**)
Wiens, Jonathan P. (*see* **Mooi, Randall D.**)
Willson, Robert J. (*see* **Rouse, Jeremy D.**)
Winterbottom, Richard (*see* **Courtenay, Walter R., Jr.**)
Wojdak, Jeremy M. (*see* **Chambers, David L.**)
Wooten, Victor G. (*see* **Leaf, Robert T.**)
Worthy, Trevor H., Alan J. D. Tennyson, Suzanne J. Hand, Henk Godthelp, and R. Paul Scofield. Terrestrial Turtle Fossils from New Zealand Refloat Moa's Ark 72
Wosiacki, Wolmar B. (*see* **Mendonça, Marina B.**)
Yarnell, Sarah M. (*see* **Kupferberg, Sarah J.**)
Zappalorti, Robert T. (*see* **Reinert, Howard K.**)
Zhou, Wei, Xu Li, and Alfred W. Thomson. A New Genus of Glyptosternine Catfish (Siluriformes: Sisoridae) with Descriptions of Two New Species from Yunnan, China 226